

**THE INFLUNCE OF DIABETES MEDICATIONS ON THE ASSOCIATIONS
BETWEEN BMI AND CARDIOVASCULAR HEALTH RISKS**

ANDREW SZIVOS

A THESIS SUBMITTED TO THE FACULTY OF GRADUATE STUDIES IN PARTIAL
FULLFILLMENT OF THE REQUIREMENTS FOR THE DEGREE OF

MASTER OF SCIENCE IN KINESIOLOGY AND HEALTH SCIENCE

YORK UNIVERSITY

TORONTO, CANADA

AUGUST 2022

©ANDREW SZIVOS, 2022

Abstract

The purpose was to observe the influence of diabetes medications on the association between body mass index (BMI) and cardiovascular health (CVH). Individuals with type 2 diabetes (T2D) from the National Health and Nutrition Examination Survey (NHANES Continuous 1999-2018) were grouped by diabetes medications with similar body weight side effects. There was a positive association between BMI for triglycerides in the combination medications group ($P<0.002$). Conversely, there were negative associations between BMI for HbA1c in the weight gain medications group ($P=0.02$), and total cholesterol in the weight loss and weight gain medication groups ($P<0.05$). Finally, no associations were between BMI in any medication group for fasting glucose, blood pressure or high-density lipoproteins ($P>0.05$). Therefore, T2D medications may weaken or invert the commonly observed association between obesity and CVH. Thus, patients living with T2D and obesity may have better CVH compared to patients without obesity taking the same T2D medication.

Acknowledgement

I would first like to thank my parents for supporting and encouraging me on this long journey. You both were there for me through the hardships of my university studies and made sure I would stay focused on what is important. My sister, who would tell me the truth even when I didn't want to hear it. Thank you for being there for me when I needed you most and motivating me when I over exaggerated my problems. I hope I was able to make you all proud of the person I have become and the path I chose to pursue.

I would also like to thank my lab mates: Rubin, Zahava, Maral, Baara, Amir and Yasmine. You all made the past 2 years of navigating between virtual and in person learning more enjoyable. Communicating over Zoom was definitely not an easy adjustment to make for anyone, but being surrounded by supportive lab mates kept me motivated to continue pressing forward and see this to the end. Thank you all for being such kind, understanding and accepting individuals as well as sharing your time with me when I needed assistance.

Finally, it is with pleasure that I thank Dr. Jennifer Kuk for taking me under her wing as a graduate student. I never could have imagined that the brief encounter we shared would lead to an entire master degree. There are not enough words to express how grateful I am for all the patience, guidance and understanding you gave me. You pushed me forward to grow, not just as a student, but as a confident individual that wants to succeed in the work I am passionate for. The time you shared with me as a graduate student will not be forgotten anytime soon as I am sure to still see you from time to time. Thank you for being strict with me when I would make the same mistakes and never giving up on me. I could not have asked for a better mentor.

Table of Contents

Abstract.....	ii
Acknowledgement.....	iii
Table of Contents	iv
List of Tables	v
List of Figure	vi
List of Abbreviations	vii
1.0 General Introduction.....	1
2.0 Review of Literature	2
2.1 Introduction.....	2
2.2 Type 2 Diabetes (T2D)	2
2.3 Managing Blood Glucose and HbA1c	2
2.4 Obesity	3
2.5 Obesity and Cardiovascular Health Risks.....	3
2.6 Medications Available for Treatment of T2D.....	4
3.0 Manuscript.....	9
3.1 Abstract.....	9
3.2 Introduction.....	9
3.3 Methods.....	10
3.4 Results.....	13
3.5 Discussion.....	19
4.0 General Conclusion.....	23
5.0 References.....	24

List of Tables

Table 1: Descriptive characteristics of US adults living with diabetes by medication class

predicted weight effects (NHANES 1999- 2018) 16

List of Figure

Figure 1: Weighted means for HbA1c, Fasting Glucose, SBP and DBP by BMI and Medication

Group.....17

Figure 2: Weighted means for Total Cholesterol, HDL and Triglyceride by BMI and Medication

Group 18

List of Abbreviations

Abbreviation	Term
BMI	Body Mass Index
CombMed	Combination Medications
DBP	Diastolic Blood Pressure
DPP	Dipeptidyl peptidase
GLP	Glucagon-like peptide
HbA1c	Hemoglobin A1c
HDL	High Density Lipoproteins
NHANES	National Health and Nutrition Examination Survey
nOB	No Obesity
OB	Obesity
OGTT	Oral Glucose Tolerance Test
SBP	Systolic Blood Pressure
SGLT	Sodium Glucose Co-Transporter
T2D	Type 2 Diabetes
TZD	Thiazolidinediones

U.S.	United States
WgMed	Weight Gain Medications
WlMed	Weight Loss Medications
WnMed	Weight Neutral Medications

1.0 General Introduction

The prevalence of obesity in United States (U.S.) adults has escalated in the past few decades from 15% in 1980 (1) to 42% in 2017 (2). Obesity is well known to be associated with developing further health complications, including type 2 diabetes (T2D) (3–5). T2D is a major chronic disease, which as of 2018 is prevalent in over 34 million U.S. adults, affecting an estimated 13% of the U.S. population (6). In 2017, 25% of all U.S. healthcare expenditures went towards diabetes care (7). Those living with T2D are advised to lower their blood glucose and glycated hemoglobin (HbA1c) levels through lifestyle management, along with antidiabetic medications (8).

Medications are effective interventions for lowering HbA1c and blood glucose levels (9). Diabetes medication work through several different physiological mechanisms and can have side effects that result in body weight gain or loss (10). However, it is unknown if diabetes medications alter the known association between BMI and other cardiovascular risk factors. Thus, the purpose of this study was to examine the potential influence diabetes medications have on the relationship between BMI and cardiovascular health risks.

2.0 Review of Literature

2.1 Introduction

Obesity is a condition that increases the risk of developing further health complications (5). One of the major health complications associated with obesity is T2D which is characterized by hyperglycemia as a result of complete or partial insufficiency of insulin secretion and/or action (11,12). T2D is the most prevalent type of diabetes accounting for 8.5% of US adult population (13).

This literature review will discuss T2D, the importance of regulating blood glucose and HbA1c, obesity and the cardiovascular health risks associated with obesity. This review will also outline the medications currently available for T2D patients and considerations when prescribing medications to individuals living with obesity.

2.2 Type 2 Diabetes (T2D)

T2D is characterized by the inability to control blood glucose levels due to beta cell dysfunction and/or insulin resistance (14). T2D is diagnosed as: an HbA1c $\geq 6.5\%$, a fasting glucose ≥ 7 mmol/L, a randomized blood glucose test ≥ 11.1 mmol/L or a two hour oral glucose tolerance test (OGTT) ≥ 11.1 mmol/L (11). Early symptoms for T2D include excessive urination, excessive thirst, increased hunger, and fatigue (15). Longer term complications can include CVD, vision changes, yeast infections, slow healing sores, and feelings of numbness in peripheral body parts (16).

2.3 Managing Blood Glucose and HbA1c

Individuals with T2D usually have elevated blood glucose levels which is known as hyperglycemia. Hyperglycemia can lead to damage of capillary endothelial cells in the retina,

mesangial cells in the renal glomerulus, and Schwann cells in peripheral nerves (17). Blood glucose can be difficult to regulate, however regular self-monitoring with the use of a glucometer makes self-management easier for the individual. HbA1c or glycated hemoglobin is a molecule formed when glucose binds with the proteins on red blood cells (18). Higher blood glucose levels increase the likelihood for glucose to bind with these proteins. Testing HbA1c is the gold standard when looking at T2D blood glucose control and indicates blood glucose levels over a 2-3 month period (19).

2.4 Obesity

Obesity is defined as an excessive amount of body fat that has negative implications on one's health (20). The most common measure used to diagnose obesity is body mass index (BMI). BMI is calculated by dividing an individual's body weight in kilograms from their height in meters squared (kg/m^2). It has been well established that a $\text{BMI} \geq 30 \text{ kg}/\text{m}^2$ indicates a person is living with obesity (4). Health quality of life such as overall physical functioning and mobility is greatly increased with reductions in body weight (22).

2.5 Obesity and Cardiovascular Health Risks

Individuals living with obesity are shown to have elevated health risks compared to those without obesity (23). It is well established that blood glucose and HbA1c levels are more likely to be elevated for individuals living with obesity as they are at increased risk for insulin resistance (24). Obesity is also associated with hypertension which is defined as blood pressure measures $\geq 140/90 \text{ mmHg}$ (25) and dyslipidemia or abnormally high cholesterol levels which is defined as triglycerides $\geq 150\text{mg}/\text{dl}$ or $\text{HDL} < 40 \text{ mg}/\text{dl}$ (26–28).

2.6 Medications Available for Treatment of T2D

Antidiabetic medications in conjunction with lifestyle are a vital part of treatment for individuals with diabetes in order to control glucose levels (9). These medications work through a number of pathways and have varied effects on body weight. Amylin analogs, and insulin are currently only effective as subcutaneous injections as they are destroyed by gastric acid before reaching the bloodstream (29,30). Glucagon-like peptide (GLP)-1-agonists are also only injectable however recent discoveries in oral options effectively lower blood glucose (31). Biguanides, sodium glucose co-transporters type 2 (SGLT2) inhibitors, alpha-glucosidase inhibitors, dipeptidyl peptidase (DPP)-4-inhibitors, sulfonylureas, meglitinides, and thiazolidinediones are all oral medication options currently available to lower blood glucose (10,32). The medications examined in this paper include the following: amylin analogs, biguanides, GLP-1-agonists, SGLT2 inhibitors, alpha-glucosidase inhibitors, DPP-4-inhibitors, insulin, sulfonylureas, meglitinides, and thiazolidinediones.

Medications Associated with Weight Loss

The following T2D medications are associated with a decrease in body weight: amylin analogs, biguanides, GLP-1-agonists and sodium glucose co-transporters type 2 (SGLT2) inhibitors (10,33–38).

Amylin Analogs

The only available amylin medication is known as pramlintide which is a synthetic compound of the hormone amylin (10). Human amylin is co-secreted with insulin after eating a meal which lowers glucose in the bloodstream (39). Pramlintide and amylin lower blood glucose by suppressing postprandial production of glucose and slowing the rate of gastric emptying

(10,33). Currently the mechanism behind weight loss is unknown but could be attributed to reductions in food intake as feelings of satiety after a meal are increased (40).

Biguanides

The most commonly used medication for T2D is metformin as it is associated with minimal negative side effects and potential improvements to cardiovascular health (11). Metformin reduces blood glucose concentration by lowering glucose production from the liver and increasing insulin sensitivity (35). The mechanism causing weight loss with metformin is thought to be associated with decreased appetite (41).

GLP-1-agonists

GLP-1 is a gut hormone which is secreted by the intestines after eating as a meal in response to increased glucose levels (42). GLP-1-agonists are medications which bind to G-protein receptors similarly to the gut hormone GLP-1. These medications lower blood glucose by increasing insulin secretion from the pancreas and inhibiting glucose production from the liver (14,34,42). Reduction in body weight may be associated with decreased gastric emptying and increases feelings of satiety (43).

SGLT2 Inhibitors

SGLT2 inhibitors lower blood glucose concentrations by increasing urinary glucose elimination and reducing renal glucose re-absorption after a meal. (44). Long term use of dapagliflozin is reported to be associated with statistically significant weight loss through reductions of visceral adipose tissue and subcutaneous adipose tissue (45). Another study demonstrates that empagliflozin is associated with a shift usage from sugars to fats for energy which may explain the reduced body weight over time (46).

Medications Associated with Neutral Changes in Body Weight

The following medications are reported to be weight neutral: alpha-glucosidase inhibitors and Dipeptidyl peptidase (DPP)-4-inhibitors (10,47,48).

Alpha-glucosidase Inhibitors

Alpha glucosidase inhibitors prevent the enzyme alpha-glucosidase from breaking down complex sugars into simple sugars (12,49). By inhibiting alpha glucosidase, the small intestine is limited to the absorption of complex sugars which lowers the rise in blood glucose levels after eating.

DPP-4-inhibitors

DPP-4-inhibitors extend the half-life of endogenous and exogenous GLP-1 by reducing the actions of DPP-4 enzymes (50). Extending the actions of GLP-1 lower blood glucose levels through the prolonged release of insulin. The use of DPP-4 inhibitors are weight neutral compared to GLP-1-agonists, indicating lower GLP-1 receptor stimulation or lower release of GLP-1 (50,51).

Medications Associated with Weight Gain

The following medications are associated with a clinically significant increase in body weight (i.e. >5-10%): insulin, sulfonylureas, meglitinides, and thiazolidinediones (TZD) (10,52–56).

Insulin

Exogenous insulin is recommended for T2D patients when oral antidiabetic agents fail to manage blood glucose levels (57). Insulin suppresses hepatic glucose production and increases

glucose uptake by adipose and muscle tissues to decrease blood glucose levels (34). Higher levels of insulin increases stimulation of lipogenesis which could result in weight gain (58).

Sulfonylureas and Meglitinides

Sulfonylureas and meglitinides both induce closure of the potassium ATP channels in beta cells which promotes insulin secretion (53,59). Promoting short term rapid insulin secretion aids in the absorption of glucose after a meal therefore lowering blood glucose. Meglitinides have a shorter half-life which is why they are less likely to cause hypoglycemia compared to sulfonylureas. (60,61). As seen with patients on exogenous insulin therapy, individuals on these medications also gain weight.

TZDs

TZDs lower blood glucose level by increasing insulin sensitivity in skeletal muscle (55). Use of TZDs are associated with weight gain resulting from accumulations of subcutaneous fat (62). These medications may also cause fluid retention (i.e. body water) in peripheral tissues which can lead to swelling and weight gain (55).

Antidiabetic Combinations

Combination medications are used as a progressive measure when individuals are unable to maintain adequate blood glucose control with a single medication. When prescribing oral combinations, one medication generally increase insulin sensitivity while the other increase secretion of insulin (63). Metformin and sulfonylureas are usually the first medications packaged followed by thiazolidinediones or alpha-glucosidase inhibitors (64,65).

Summary of Literature

It is well known that obesity is associated with higher health risk along with increased risk for the development of T2D. Those living with T2D may be prescribed medications with weight altering effects which may change the commonly observed associations between obesity and cardiovascular health risks. The aim of this study is to examine if diabetes medications alter the associations between BMI and CVH.

3.0 Manuscript

3.1 Abstract

The purpose was to observe the influence of diabetes medications on the association between body mass index (BMI) and cardiovascular health (CVH). Individuals with type 2 diabetes (T2D) from the National Health and Nutrition Examination Survey (NHANES Continuous 1999-2018) were grouped by diabetes medications with similar body weight side effects. There was a positive association between BMI for triglycerides in the combination medications group ($P < 0.002$). Conversely, there were negative associations between BMI for HbA1c in the weight gain medications group ($P = 0.02$), and total cholesterol in the weight loss and weight gain medication groups ($P < 0.05$). Finally, no associations were between BMI in any medication group for fasting glucose, blood pressure or high-density lipoproteins ($P > 0.05$). Therefore, T2D medications may weaken or invert the commonly observed association between obesity and CVH. Thus, patients living with T2D and obesity may have better CVH compared to patients without obesity taking the same T2D medication.

3.2 Introduction

Obesity and T2D both lead to severe health outcomes. The use of various anti-hyperglycemic medications are effective in lowering blood glucose levels though different mechanisms, with some promoting weight gain or weight loss. However, the effects these medications for T2D have on the health of individuals independent of BMI is still not fully understood. Therefore, the objective of this study is to examine the influence anti-hyperglycemic medications may have on the associations between BMI and cardiovascular health risk.

3.3 Methods

Participants

The data in this study was collected by the National Health and Nutrition Examination Survey (NHANES) Continuous 1999 - 2018 (n= 101 316). NHANES is a nationally represented survey in the United States conducted every 2 years focusing on a wide range of different health topics (66). The survey data for NHANES is collected through home interviews, followed by a health examination and laboratory samples which are conducted in mobile medical examination centers (MEC). In the home interviews, participants are asked demographic, socioeconomic and health related questions. During the health examinations, medical, dental and physiological measurements are taken. Furthermore, laboratory samples are taken by trained medical personnel. All participants gave written informed consent before participating.

Participants for this study were included if they had diabetes as defined by the following criteria: were told by a doctor they have diabetes (n= 6807), fasting glucose ≥ 7 mmol/L (n= 2587), random blood glucose ≥ 11.1 mmol/L (n= 783), HbA1c level $\geq 6.5\%$ (n = 5578), currently on a diabetes prescription medication (n = 6011), or 2-hour glucose tolerance test ≥ 11.1 mmol/L (n = 810). Individuals were excluded if they had missing data for body weight (n= 6134), height (n= 13 077), or HbA1c (n= 36 364). Individuals were also excluded if they were only taking insulin (n= 517), age < 20 years (n= 46 235), or answered “yes” to the question “Are you pregnant?” (n= 1722). An analytical sample of 7953 were used for the final analysis. Outliers for each cardiovascular health risk were excluded for each relevant analysis: (HbA1c n = 3, SBP n = 1, DBP n = 1, total cholesterol n = 4, HDL n = 2, Triglyceride n = 4). BMI was stratified into two groups: no obesity (nOB, BMI < 30 kg/m²) and obesity (OB, BMI ≥ 30 kg/m²).

Interview and Examination Measures

Demographic information was collected through home interviews and questionnaires prior to the health examination in the MEC. Sex was recorded as male or female. Individuals older than 85 years of age were recoded as 85 years of age for disclosure reasons. Ethnicity was recoded as either Non-Hispanic White or Other Race/Multi-Racial. All body measurements were conducted by trained professional in the MEC. Body weight was measured using a digital scale. Standing height was measured using a stadiometer with a fixed backboard and an adjustable head piece. BMI was calculated as body weight (kg) and divided by height squared (m^2) (67). Waist circumference was measured in centimeters at the level of the uppermost lateral border of the right ilium (68).

For smoking status, individuals were asked if they smoked at least 100 cigarettes in their lifetime. If they answered “no” they were classified as nonsmokers. If they answered “yes” they were then asked how often they currently smoke cigarettes in a month. If participants answered “every day” or “some days” they were classified as current smokers. If participants answered “not at all”, “refused”, “don’t know” or had missing data, they were classified as a past smoker.

Socioeconomic status of participants was determined using the poverty income ratio (PIR). The Department of Health and Human Services’ (HHS) poverty guidelines were used as the poverty measure to calculate this index. The values were calculated by dividing the family income by the poverty guidelines for the specific amount of family members set by year and state. Individuals who had values greater than 5 were recoded as 5 for disclosure concerns. A mean substitution was also performed for individuals who had missing data for this variable. If participants were lower than a value of 1, they were considered under the poverty threshold (69).

Diabetes Medication Use

NHANES uses the Lexicon Plus database which contains medication ingredients, medication classifications and medication uses. There are currently 11 classes of antidiabetic agents which can be prescribed for diabetes: alpha-glucosidase inhibitors, amylin analogs, antidiabetic combinations, biguanides, dpp-4-inhibitors, glp-1-agonists, insulin, meglitinides, sglt-2-inhibitors, sulfonylureas and thiazolidinediones. A total of 5 medication groups (Medgroups) were created grouped by body weight side effects. The no medication (NoMed) group consisted of individuals currently not on diabetes medications but still classified as a person living with diabetes. The weight loss medication (WlMed) group consisted of individuals taking diabetes medications associated with weight loss (10,36,38,70,71): biguanides, amylin analogs, glp-1-agonists, and sglt-2-inhibitors. The weight neutral medication (WnMed) group consisted of individuals taking diabetes medications that are not reported to have an effect on weight (10,72,73): alpha-glucosidase inhibitors, and dpp-4-inhibitors. The weight gain medication (WgMed) group consisted of individuals taking diabetes medication associated with weight gain (10,54,74,75): sulfonylureas, insulin, meglitinides or thiazolidinediones. Finally, the combination medication (CombMed) group consisted of individuals on more than one diabetes medication or taking antidiabetic combination medications.

Statistical Analysis

Descriptive characteristics were stratified by diabetes medication weight side effect groups. Data was shown as means (SE) or prevalence % (SE). MEC weighting variable was used for HbA1c, SBP, and DBP analyses while the fasting (SAF) weighting variable was used for analyses with fasting glucose, total cholesterol, HDL and triglyceride. Linear regression analysis was used to assess the association between HbA1c, SBP, DBP, fasting glucose, total cholesterol

and HDL and triglycerides with BMI (continuous variable) and medication groups, with examination of main effects and interactions. With significant BMI-med group main effects and/or interactions, the regression was repeated with BMI was coded as nOB and OB (BMI cut-off 30 kg/m²) to facilitated graphical representation and least squared means post hoc tests were used to determine obesity-medication group differences. All means were adjusted for sex, ethnicity, age, socioeconomic status, and smoking status. SBP and DBP analysis were further adjusted for high blood pressure medications. Total cholesterol, high density lipoproteins (HDL), and triglyceride analysis were further adjusted for lipid medications.

All analyses were performed using SAS version 9.4 and weighted to be nationally representative of the US population (SAS Institute Inc., Cary, NC, USA). Results were statistically significant if $P < 0.05$.

3.4 Results

Descriptive Characteristics

Participant characteristics are shown in **Table 1**. The analysis consisted of 5 (medgroups) stratified based on body weight side effects: (NoMed n= 3161), (WlMed n=1453), (WnMed n= 47), (WgMed n= 756), and (CombMed n= 2536). Individuals in the NoMed group were on average younger (57.5 ± 0.4 years) than those in any other medgroup (58.4-68.0 years). The average fasting glucose in the CombMed group was the highest (9.2 ± 0.2 mmol/L) compared to all the other medgroups (7.3-8.1 mmol/L; $P < .0001$).

Association between BMI and Diabetic Medication use in Cardiovascular Health Risks

The weighted means for HbA1c, fasting glucose, SBP and DBP stratified by BMI group and medgroup are shown in **Figure 1**. Within the whole sample, there was no simple correlation between BMI and HbA1c ($P=0.76$). For HbA1c, the interaction effect between BMI and

medgroup trended towards significance ($P=0.08$) with a significant main effect of medgroup ($P < 0.0001$). When stratified by medication group, there was a negative association between BMI and HbA1c in the WgMed group ($P= 0.02$), with no significant associations between BMI and HbA1c in the other medgroups. The HbA1c in the CombMed group ($nOB: 7.9 \pm 0.1\%$; $OB: 7.7 \pm 0.1\%$) was significantly greater than all medgroups for both BMI groups ($nOB: 6.7-7.2\%$, $OB: 6.7-7.0\%$, $P < 0.05$). Furthermore, within those with obesity, the HbA1c in the WlMed group was lower than all other medgroups ($6.7 \pm 0.1\%$ versus $6.8- 7.7$, $P < 0.05$).

Within the whole sample, there was no simple correlation between BMI and fasting glucose ($P= 0.83$). For fasting glucose, the interaction effect between BMI and medgroups trended towards significance ($P=0.09$) with a significant main effect of medgroup ($P= 0.0023$). When stratified by medication group, there was a significant positive association between BMI and fasting glucose in the NoMed group ($P= 0.02$) with no significant associations between BMI and fasting glucose in the other medgroups. Individuals in the CombMed group had significantly higher fasting glucose levels for a given BMI group ($nOB: 9.2 \pm 0.3$ mmol/L; $OB: 9.2 \pm 0.2$ mmol/L) compared to the NoMed, WlMed, and WgMed groups ($nOB: 7.3- 8.3$ mmol/L; $OB: 7.8- 8.2$ mmol/L; $P < 0.05$). The interaction effect between BMI and medgroup along with both main effects of BMI and medgroup were not significant for SBP and DBP ($P > 0.05$).

The weighted means for total cholesterol, HDL, and TG are shown in **Figure 2**. There was a simple negative correlation between BMI and total cholesterol within the whole sample ($P= 0.04$). For total cholesterol, the interaction effect between BMI and medgroups trended towards significance ($P= 0.07$), with a significant main effect of medgroup ($P= 0.0061$). When stratified by medication group and adjusted for demographic variables, socioeconomic status,

smoking status and lipid medications, the association remained negative between BMI and total cholesterol in the NoMed, WlMed and WgMed groups ($P < 0.05$), but with no significant associations between BMI and total cholesterol in the WnMed or CombMed groups. Furthermore, total cholesterol in the NoMed group was significantly higher compared to the WlMed and CombMed group for both BMI groups ($P < 0.05$).

Within the whole sample, there was no simple correlation between BMI and TG ($P = 0.28$). For TG, the interaction effect between BMI and medgroup was significant ($P = 0.02$), with a significant main effect of medgroup ($P = 0.002$). When stratified by medication group and adjusted for demographic variables, socioeconomic status, smoking status and lipid medications, there was a positive association between BMI and TG in the CombMed group ($P = 0.002$), but with no significant associations between BMI and TG in the other medgroups. For HDL, there was a simple negative correlation for BMI ($P < 0.0001$), with no significant interaction effect between BMI and medgroup or main effect of medgroup ($P > 0.05$).

Table 1: Descriptive characteristics of US adults living with diabetes by medication class predicted weight effects (NHANES 1999- 2018)

Participant Characteristics	No Medication (n= 3161)	Weight Loss Medication (n=1453)	Weight Neutral Medication (n= 47)	Weight Gain Medication (n= 756)	Combination Medication (n= 2536)
Variable	Mean (SE)/ Freq, SE	Mean (SE)/ Freq, SE	Mean (SE)/ Freq, SE	Mean (SE)/ Freq, SE	Mean (SE)/ Freq, SE
Age (years)	57.5 (0.4) ^{cde}	58.4 (0.5) ^{cde}	68.0 (2.1) ^{abe}	65.2 (0.7) ^{abe}	60.6 (0.4) ^{abcd}
Sex (% female)	51.0, 0.1 ^b	45.9, 0.1 ^{ae}	49.9, 0.1	50.4, 0.1	53.8, 0.1 ^b
Ethnicity (% white)	49.0, 0.1 ^{bde}	56.2, 0.1 ^{ad}	38.3, 0.1	38.0, 0.1 ^{abe}	54.2, 0.1 ^{ad}
Smoking Status (% smokers)	19.7, 0.1 ^{bde}	13.1, 0.1 ^a	10.4, 0.1	13.0, 0.1 ^a	14.7, 0.1 ^a
PIR (% below poverty)	16.2, 0.1 ^{bd}	12.6, 0.1 ^a	10.8, 0.1	11.9, 0.1 ^a	14.8, 0.1
Body Weight (kg)	90.8 (0.6) ^{de}	93.1 (1.0) ^d	86.9 (3.6) ^e	88.5 (1.0) ^{abe}	95.1 (0.8) ^{acd}
Height (cm)	167.4 (0.3) ^d	166.6 (0.4) ^e	167.9 (2.8)	165.5 (0.5) ^{ae}	168.2 (0.3) ^{bd}
BMI (kg/m ²)	32.3 (0.2) ^{be}	33.4 (0.3) ^{ad}	31.1 (1.3)	32.2 (0.3) ^{be}	33.5 (0.2) ^{ad}
Waist Circumference (cm)	108.7 (0.4) ^{be}	111.1 (0.6) ^{ad}	107.5 (2.2) ^e	109.1 (0.7) ^{be}	112.7 (0.6) ^{acd}
Cardiovascular Health Risks					
Fasting Glucose (mmol/L)	7.8 (0.1) ^e	7.9 (0.1) ^e	7.3 (0.4) ^e	8.1 (0.2) ^e	9.2 (0.2) ^{abcd}
HbA1c (%)	6.8 (0.1) ^e	6.8 (0.1) ^{de}	7.3 (0.5)	7.0 (0.1) ^{be}	7.7 (0.1) ^{abd}
Systolic Blood Pressure (mmHg)	131 (1) ^{bd}	129 (1) ^{ade}	133 (4)	136 (1) ^{abe}	131 (1) ^{bd}
Diastolic Blood Pressure (mmHg)	72 (1) ^{cde}	71 (1) ^{cde}	66 (2) ^{ab}	69 (1) ^{ab}	68 (1) ^{ab}
Total Cholesterol (mmol/L)	5.2 (0.1) ^{bcd}	4.7 (0.1) ^a	4.5 (0.3) ^a	4.8 (0.1) ^{ae}	4.6 (0.1) ^{ad}
HDL (mmol/L)	1.3 (0.1) ^e	1.3 (0.1) ^e	1.3 (0.1)	1.3 (0.1) ^e	1.2 (0.1) ^{abd}
Triglyceride (mmol/L)	2.0 (0.1) ^c	1.9 (0.1)	1.5 (0.3) ^a	2.1 (0.2)	2.0 (0.1)

Weighted sample size: Medical Examination Centre = 298,416,774, Sample after fasting = 260,805,647, Waist Circumference (n) = 7555 Fasting Glucose (n) = 3936 SBP (n) = 7638 DBP (n) = 7592 Total Cholesterol (n) = 3885 HDL (n) = 3885 Triglyceride (n) = 3369

a = Significantly different compared to the no medications group (P < 0.05), b = Significantly different compared to the weight loss medication group (P < 0.05), c = Significantly different compared to the weight neutral medication group (P < 0.05), d = Significantly different compared to the weight gain medication group (P < 0.05), e = Significantly different compared to the combination medication group (P < 0.05)




Figure 1: Weighted means for HbA1c: $n = 7644$, Fasting Glucose: $n = 3631$, SBP: $n = 7359$, and DBP $n = 7295$ by BMI and Medication Groups. Means are adjusted for sex, ethnicity, age, social economic status and smoking status. SBP and DBP are further adjusted for high BP medication status. Medgroup= medication group, nOB= BMI <30 (kg/m^2), OB= BMI ≥ 30 (kg/m^2), BMI= body mass index, SBP= systolic blood pressure, DBP= diastolic blood pressure.

*= Significant BMI difference within medication group ($P < 0.05$). a= Significantly different from NoMed group within BMI group, b= Significantly different from WMed group within BMI group, c= Significantly different from WnMed group within BMI group, d= Significantly different from WgMed group within BMI group, e= Significantly different from CombMed group within BMI group.

NoMed= No Medications, WMed= biguanides, amylin analogs, glp-1-agonists, and sglt-2-inhibitors, WnMed= alpha-glucosidase inhibitors, and dpp-4-inhibitors, WgMed= sulfonylureas, insulin, meglitinides or thiazolidinediones, CombMed= more than one medication and antidiabetic combination medications.




Figure 2: Weighted means for Total Cholesterol: $n = 3579$, HDL: $n = 3579$, and Triglyceride: $n = 3111$ by BMI and Medication Groups. Means are adjusted for sex, ethnicity, age, social economic status and smoking status and lipid medication status. Medgroup= medication group, nOB= BMI < 30 (kg/m^2), OB= BMI ≥ 30 (kg/m^2), BMI= body mass index, HDL= high density lipoproteins.

*= Significant BMI difference within medication group ($P < 0.05$), a= Significantly different from NoMed group within BMI group, b= Significantly different from WlMed group within BMI group, c= Significantly different from WnMed group within BMI group, d= Significantly different from WgMed group within BMI group, e= Significantly different from CombMed group within BMI group.

NoMed= No Medications, WlMed= biguanides, amylin analogs, glp-1-agonists, and sgl-2-inhibitors, WnMed= alpha-glucosidase inhibitors, and dpp-4-inhibitors, WgMed= sulfonylureas, insulin, meglitinides or thiazolidinediones, CombMed= more than one medication and antidiabetic combination medications.

3.5 Discussion

Several medications used to treat hyperglycemia in T2D promote either weight loss or weight gain as a side effect of treatment. To our knowledge, the current study is the first to demonstrate that certain diabetes-related medications that lower glycemia but also impact body weight (increase or decrease) can potentially alter the commonly reported positive association between BMI with glycemic control and total cholesterol levels. Specifically, we found that those with T2D and obesity taking WIMeds were associated with lower HbA1c levels compared to those with T2D without obesity taking the same medication. As well, those with T2D and obesity taking WIMeds or WgMeds were associated with lower total cholesterol levels compared to those with T2D without obesity taking the same medications. Thus, diabetes medications may alter how obesity relates with cardiometabolic risk.

Excess body fat often leads to impaired whole body insulin sensitivity, beta-cell dysfunction and dysglycemia (14,76). Thus, those with a higher BMI are more likely to have T2D (77,78). T2D medications lower glucose levels by numerous mechanisms with some promoting clinically meaningful weight gain (i.e. insulin) while others promoting clinically meaningful weight loss (i.e., most oral hypoglycemic agents) (9). Thus, the known positive association between BMI and HbA1c may change when taking certain diabetes medications. Within the full sample of adults with T2D, there was no simple correlation between BMI and HbA1c observed in our study. In fact, those living with obesity and taking diabetes WIMeds had the lowest mean HbA1c ($6.7 \pm 0.1\%$) of all obesity-medication groups. Indeed, certain diabetes medications may improve glucose control and improve glucose utilization through various mechanisms that may or may not be independent of obesity. One of the effects of WIMeds such as amylin analogs, metformin and GLP-1-agonists is loss of appetite or feelings of satiety

(40,41,43). As such, individuals may lower HbA1c levels through reductions in obesity. Conversely, certain diabetes medications are reported to be associated with improvements in HbA1c despite concurrent weight gain (79,80). We observed a negative association between BMI and HbA1c for those taking diabetes WgMeds. The glucose lowering mechanism for diabetes WgMeds such as sulfonylureas, TZDs and meglitinides is increased insulin production (34,55,59). Greater insulin leads to lower blood glucose, greater energy storage, and weight gain as a consequence. Thus, while the association between BMI and HbA1c is often positive, this study demonstrates that diabetes medications that alter body weight may also alter and even invert the association between BMI and HbA1c.

It is commonly reported that individuals with obesity have higher total cholesterol levels compared to individuals without obesity (81,82). In contrast, we observe a simple negative association between BMI and total cholesterol in our population of adults with T2D. This negative association remained true for those taking NoMeds, WlMeds or WgMeds. Cholesterol synthesis is known to be positively associated with increased adipose tissues which stores cholesterol as triglyceride for later use (83,84). Many diabetes medications increase insulin production and sensitivity (10), which can increase the storage of triglycerides and lower total cholesterol. Thus, diabetes medications may be associated with improvements in total cholesterol despite increased BMI. We observed individuals taking WlMeds or CombMeds in our study had lower total cholesterol values than those on NoMeds for a given BMI. It is known that WlMeds, notably metformin may also reduce total cholesterol, specifically LDL cholesterol (85), potentially due to decreased appetite and reduced food intake (40,41,43). Together, these results suggest that certain factors such as T2D medications can significantly alter the often observed positive association between BMI and total cholesterol.

In this study, we demonstrate that the association between obesity and health risk in those taking diabetes medications may be altered. This may be expected given the several different mechanisms of action possible for diabetes medications. Although the association between obesity and health risk is well established (86,87), there are several other factors that are known to alter this association such as aging, physical activity and diet (88–90). Unlike past studies, BMI was not significantly associated with the majority of cardiovascular health risk factors examined in the current analyses. There are studies which suggest that the association between BMI and cardiovascular health risk is weaker in older individuals (91). In fact, some report inversed associations between BMI and mortality risk in older adults (92,93). These differences with aging, may have contributed to the unexpectedly weak association between obesity and health risk in this study as individual with T2D are generally older. Similarly, it is known that there are chronic and acute effects of physical activity which may obscure the association between BMI and CVH (94,95). Those living with T2D are advised at least 150 minutes a week of moderate to vigorous aerobic physical activity (96), but only 25% of individuals meet this guideline (97). Similarly, diet is known to have acute and chronic effects on cardiometabolic health risk (98,99). Although patients with T2D are prescribed a Mediterranean style diet (100,101), the adherence to this recommended dietary approach is low with less than 50% of individuals complying (102). Further, the association between dietary factors and health is complicated and often hard to capture with a single measure. Therefore, there are several factors along with diabetes medications that may influence the association between BMI and CVH.

There were strengths and limitations to this study. The use of NHANES allowed for easy access over a long period of time with a large sample size weighted to be nationally representative of the US population (66). Furthermore, the use of home interviews lowered the

amount of nonresponse for demographic data and allowed for confounding adjustments in our analysis. The main limitation to this study was the use of cross sectional data which prevents the study from examining causal relationships. The sample size for individuals using WnMeds (n= 47) was small compared to other Medgroups thereby limiting our ability to observe significant group differences. Lastly, the duration or dosage in which individuals were taking specific diabetes medications was not assessed.

In conclusion, our study showed that those with T2D and obesity on WlMeds were associated with lower HbA1c levels compared to those with T2D without obesity on the same medication. Furthermore, those with T2D and obesity on WlMeds and WgMeds were associated with lower total cholesterol levels compared to those with T2D without obesity on the same medication. These findings imply the relationship between obesity status and cardiovascular health risks are complex and can change when taking diabetes medications. Further research should be conducted on those with T2D living with obesity have better long term health outcomes compared to those with T2D without obesity taking the same medication.

4.0 General Conclusion

Obesity is a rising health issue globally (1) which increases an individual's risk for developing further diseases, including T2D (3). The global prevalence of T2D is also on the rise with the number of individuals living with this disease predicted to climb over 366 million people by 2030 (103). As such, there is an increasing focus on T2D management, where lifestyle changes along with medications are crucial to reduce health outcomes.

Diabetes medications effectively reduce cardiovascular health risks such as fasting glucose and HbA1c (104), with various effects on body weight, such as weight loss or weight gain (10). There are several studies examining health changes for those living with T2D when taking diabetic medications (105,106). However, we further the literature by observing if these medications may be associated with differences in health risks between those living with and without obesity taking the same medication.

In conclusion, our study demonstrated that certain medications may be associated with greater control in HbA1c and total cholesterol levels for those living with obesity compared to without obesity when using the same medications. This suggests that in those using diabetes medications, the commonly observed positive associations between BMI and cardiovascular health risks may be altered. Further research should examine whether individuals with T2D living with obesity have better long term outcomes compared to those without obesity taking the same medication.

5.0 References

1. Freedman DS. Obesity - United States, 1988-2008. [Internet]. Vol. 60 Suppl, Morbidity and mortality weekly report MMWR. Surveillance summaries / CDC. 2011. p. 73–7.
Available from:
<https://www.cdc.gov/mmwr/preview/mmwrhtml/su6001a15.htm#:~:text=The prevalence of obesity in the United States,5%25 to 17%25 among children and adolescents%282%2C3%29.>
2. Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of Obesity and Severe Obesity Among Adults: United States, 2017-2018. NCHS Data Brief. 2020;(360):1–8.
3. Golay A, Ybarra J. Link between obesity and type 2 diabetes. Best Pract Res Clin Endocrinol Metab. 2005;19(4):649–63.
4. Hussain A, Hydrie MZ., Claussen B, Asghar S. Type 2 Diabetes and Obesity A Review. J Diabetol. 2010;2(1).
5. Kopelman PG. Obesity as a Medical Problem. 2000;404(April):635–43. Available from:
www.nature.com
6. National Diabetes Statistics Report, 2020. US Department of Health and Human Services CDC Centers for Disease Control and Prevention. 2020.
7. Yang W, Dall TM, Beronjia K, Lin J, Semilla AP, Chakrabarti R, et al. Economic costs of diabetes in the U.S. in 2017. Diabetes Care. 2018;41(5):917–28.
8. Diabetes Canada. 2018 Clinical Practice Guidelines: Quick Reference Guide. 2018.
9. He ZX, Zhou ZW, Yang Y, Yang T, Pan SY, Qiu JX, et al. Overview of clinically

- approved oral antidiabetic agents for the treatment of type 2 diabetes mellitus. *Clin Exp Pharmacol Physiol*. 2015;42(2):125–38.
10. Hermansen K, Mortensen LS. Bodyweight Changes Associated with Antihyperglycaemic Agents in Type 2 Diabetes Mellitus. *Drug Saf*. 2007;30(12):1127–42.
 11. Fox CS, Golden SH, Anderson C, Bray GA, Burke LE, De Boer IH, et al. Update on prevention of cardiovascular disease in adults with type 2 diabetes mellitus in light of recent evidence: A scientific statement from the American Heart Association and the American diabetes association. *Diabetes Care*. 2015;38(9):1777–803.
 12. Wu Y, Ding Y, Tanaka Y, Zhang W. Risk factors contributing to type 2 diabetes and recent advances in the treatment and prevention. *Int J Med Sci*. 2014;11(11):1185–200.
 13. Xu G, Liu B, Sun Y, Du Y, Snetselaar LG, Hu FB, et al. Prevalence of diagnosed type 1 and type 2 diabetes among US adults in 2016 and 2017: Population based study. *BMJ*. 2018;362.
 14. Chatterjee S, Khunti K, Davies MJ. Type 2 diabetes. *Lancet* [Internet]. 2017;389(10085):2239–51. Available from: [http://dx.doi.org/10.1016/S0140-6736\(17\)30058-2](http://dx.doi.org/10.1016/S0140-6736(17)30058-2)
 15. Diabetes [Internet]. World Health Organization. 2021. Available from: <https://www.who.int/news-room/fact-sheets/detail/diabetes>
 16. Nutrition I. Type 2 Diabetes: Symptoms, Treatments, and Long-Term Outlook [Internet]. 2021. Available from: <https://1md.org/health-guide/heart/disorders/type-2-diabetes>
 17. Brownlee M. The pathobiology of diabetic complications: A unifying mechanism.

- Diabetes. 2005;54(6):1615–25.
18. Bunn HF, Gabbay KH, Gallop PM. The glycosylation of hemoglobin: Relevance to diabetes mellitus. *Science* (80-). 1978;200(4337):21–7.
 19. Zimmet P, Alberti KG, Magliano DJ, Bennett PH. Diabetes mellitus statistics on prevalence and mortality: Facts and fallacies. *Nat Rev Endocrinol*. 2016;12(10):616–22.
 20. Wharton S, Lau DCW, Vallis M, Sharma AM, Biertho L, Campbell-Scherer D, et al. Obesity in adults: A clinical practice guideline. *Cmaj*. 2020;192(31):E875–91.
 21. Jensen MD, Ryan DH, Apovian CM, Ard JD, Comuzzie AG, Donato KA, et al. 2013 AHA/ACC/TOS Guideline for the Management of Overweight and Obesity in Adults. *J Am Coll Cardiol*. 2014;63(25):2985–3023.
 22. Fontaine KR, Barofsky I. Obesity and health- related quality of life. *Obes Rev*. 2001;2:173–82.
 23. Han TS, Sattar N, Lean M. ABC of obesity: Assessment of obesity and its clinical implications. *Br Med J*. 2006;333(7570):695–8.
 24. Modan M, Halkin H, Almog S, Lusky A, Eshkol A, Shefi M, et al. Hyperinsulinemia. A link between hypertension obesity and glucose intolerance. *J Clin Invest*. 1985;75(3):809–17.
 25. Mertens IL, Van Gaal LF. Overweight, obesity, and blood pressure: the effects of modest weight reduction. *Obes Res*. 2000;8(3):270–8.
 26. Pi-Sunyer FX. Medical hazards of obesity. *Ann Intern Med*. 1993;119(7 II):655–60.
 27. Miller WM, Nori-Janosz KE, Lillystone M, Yanez J, McCullough PA. Obesity and lipids.

- Curr Cardiol Rep. 2005;7(6):465–70.
28. G.T. G, R.S. W. Pathophysiology, Diagnosis, and Management of Dyslipidemia. Curr Probl Cardiol [Internet]. 2006;31(7):445–86. Available from:
<http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed7&NEWS=N&AN=2006311929>
 29. What is Type 1 Diabetes? [Internet]. CDC Centers for Disease Control and Prevention. 2021. Available from: <https://www.cdc.gov/diabetes/basics/what-is-type-1-diabetes.html>
 30. Traina AN, Kane MP. Primer on Pramlintide, an amylin analog. Diabetes Educ. 2011;37(3):426–31.
 31. Pechenov S, Revell J, Will S, Naylor J, Tyagi P, Patel C, et al. Development of an orally delivered GLP-1 receptor agonist through peptide engineering and drug delivery to treat chronic disease. Sci Rep [Internet]. 2021;11(1):1–15. Available from:
<https://doi.org/10.1038/s41598-021-01750-0>
 32. Mitri J, Hamdy O. Diabetes medications and body weight. Expert Opin Drug Saf. 2009;8(5):573–84.
 33. Hoogwerf BJ, Doshi KB, Diab D. Pramlintide, the synthetic analogue of amylin: Physiology, pathophysiology, and effects on glycemic control, body weight, and selected biomarkers of vascular risk. Vasc Health Risk Manag. 2008;4(2):355–62.
 34. Kurukulasuriya LR, Sowers JR. Therapies for type 2 diabetes: Lowering HbA1c and associated cardiovascular risk factors. Cardiovasc Diabetol. 2010;9:1–13.
 35. Bailey C, Path MR., Turner R. Metformin Drug Therapy. N Engl J Med.

1996;334(9):574–9.

36. Pavo I, Jermendy G, Varkonyi TT, Kerenyi Z, Gyimesi A, Shoustov S, et al. Effect of pioglitazone compared with metformin on glycemic control and indicators of insulin sensitivity in recently diagnosed patients with type 2 diabetes. *J Clin Endocrinol Metab.* 2003;88(4):1637–45.
37. Aaseth J, Ellefsen S, Alehagen U, Sundfør TM, Alexander J. Diets and drugs for weight loss and health in obesity – An update. *Biomed Pharmacother.* 2021;140(May):0–2.
38. Sohail E, Ahsan T, Ghaus S, Aijaz W. SGLT 2 inhibitors; glycemic control, weight loss and safety profile in patients with type 2 diabetes, at Medicell Institute (MIDEM). *Pakistan J Med Sci.* 2021;37(1):1–6.
39. Weyer C, Maggs D, Young A, Kolterman O. Amylin Replacement With Pramlintide as an Adjunct to Insulin Therapy in Type 1 and Type 2 Diabetes Mellitus: A Physiological Approach Toward Improved Metabolic Control. *Curr Pharm Des.* 2005;7(14):1353–73.
40. Chapman I, Parker B, Doran S, Feinle-Bisset C, Wishart J, Strobel S, et al. Effect of pramlintide on satiety and food intake in obese subjects and subjects with type 2 diabetes. *Diabetologia.* 2005;48(5):838–48.
41. Yerevanian A, Soukas A. Metformin: Mechanisms in Human Obesity and Weight Loss. *Curr Obes Rep.* 2019;8(2):156–64.
42. Ryan D, Acosta A. GLP-1 receptor agonists: Nonglycemic clinical effects in weight loss and beyond. *Obesity.* 2015;23(6):1119–29.
43. Burgmaier M, Heinrich C, Marx N. Cardiovascular effects of GLP-1 and GLP-1-based

- therapies: Implications for the cardiovascular continuum in diabetes? *Diabet Med.* 2013;30(3):289–99.
44. Bays H. Sodium glucose co-transporter type 2 (SGLT2) inhibitors: Targeting the kidney to improve glycemic control in diabetes mellitus. *Diabetes Ther.* 2013;4(2):195–220.
 45. Bolinder J, Ljunggren Ö, Kullberg J, Johansson L, Wilding J, Langkilde AM, et al. Effects of dapagliflozin on body weight, total fat mass, and regional adipose tissue distribution in patients with type 2 diabetes mellitus with inadequate glycemic control on metformin. *J Clin Endocrinol Metab.* 2012;97(3):1020–31.
 46. Brown E, Wilding JPH, Barber TM, Alam U, Cuthbertson DJ. Weight loss variability with SGLT2 inhibitors and GLP-1 receptor agonists in type 2 diabetes mellitus and obesity: Mechanistic possibilities. *Obes Rev.* 2019;20(6):816–28.
 47. LAAR FA VAN DE, LUCASSEN PL, AKKERMANS RP, LISDONK EH VAN DE, RUTTEN GE, WEEL C VAN. Alpha Glucosidase Inhibitors for Patients with Type 2 Diabetes. *Diabetes Care.* 2005;28(1):154–63.
 48. Dicker D. DPP-4 Inhibitors: Impact on glycemic control and cardiovascular risk factors. *Diabetes Care.* 2011;34(SUPPL. 2):276–8.
 49. Standl E, Schnell O. Alpha-glucosidase inhibitors 2012-cardiovascular considerations and trial evaluation. *Diabetes Vasc Dis Res.* 2012;9(3):163–9.
 50. Brunton S. GLP-1 receptor agonists vs. DPP-4 inhibitors for type 2 diabetes: Is one approach more successful or preferable than the other? *Int J Clin Pract.* 2014;68(5):557–67.

51. Thornberry N, Gallwitz B. Mechanism of action of inhibitors of dipeptidyl peptidase 4 DPP-4. *Best Pract Res Clin Endocrinology Metab.* 2009;23:479–86.
52. Goudswaard AN, Stolk RP, Zuithoff P, De Valk HW, Rutten GE. Starting insulin in type 2 diabetes: Continue oral hypoglycemic agents? A randomized trial in primary care. *J Fam Pract.* 2004;53(5):393–9.
53. Black C, Donnelly P, McIntyre L, Royle PL, Shepherd JP, Thomas S. Meglitinide analogues for type 2 diabetes mellitus. *Cochrane Database Syst Rev.* 2007;(2).
54. Sola D, Rossi L, Schianca GPC, Maffioli P, Bigliocca M, Mella R, et al. Sulfonylureas and their use in clinical practice. *Arch Med Sci.* 2015;11(4):840–8.
55. Yki-Järvinen H. Drug Therapy Thiazolidinediones. *N Engl J Med.* 2004;351(11):1106–18.
56. Diamant M, Heine RJ. Thiazolidinediones In Type 2 Diabetes Mellitus. *Drugs.* 2003;63(13):1373–405.
57. Wright A. Efficacy of addition of insulin over 6 years in patients with type 2 diabetes in the U.K. Prospective Diabetes Study. *Diabetes Care* [Internet]. 2002;25(2):330–6. Available from: <http://care.diabetesjournals.org/content/25/2/330.full.pdf+html>
58. Kersten S. Mechanisms of nutritional and hormonal regulation of lipogenesis. *EMBO Rep.* 2001;2(4):282–6.
59. Ashcroft FM. K ATP channels and islet hormone secretion : new insights and controversies. *Nat Rev Endocrinol.* 2018;9(11):660–9.
60. Varvaki Rados D, Catani Pinto L, Reck Remonti L, Bauermann Leitão C, Gross JL. The Association between Sulfonylurea Use and All-Cause and Cardiovascular Mortality: A

- Meta-Analysis with Trial Sequential Analysis of Randomized Clinical Trials. *PLoS Med.* 2016;13(4):1–22.
61. Apovian CM, Okemah J, O’Neil PM. Body Weight Considerations in the Management of Type 2 Diabetes. *Adv Ther* [Internet]. 2019;36(1):44–58. Available from: <https://doi.org/10.1007/s12325-018-0824-8>
 62. Kushner RF, Sujak M. Prevention of weight gain in adult patients with type 2 diabetes treated with pioglitazone. *Obesity.* 2009;17(5):1017–22.
 63. Nathan DM, Buse JB, Davidson MB, Heine RJ, Holman RR, Sherwin R, et al. Management of hyperglycemia in type 2 diabetes: A consensus algorithm for the initiation and adjustment of therapy. A consensus statement from the American diabetes association and the European association for the study of diabetes. *Diabetes Care.* 2006;29(8):1963–72.
 64. John M, Gopinath D, Kalra S. Triple fixed drug combinations in type 2 diabetes. *Indian J Endocrinol Metab.* 2015;19(3):311–3.
 65. Diabetes Canada Clinical Practice Guidelines. Pharmacologic Glycemic Management of Type 2 Diabetes in Adults [Internet]. 2018. Available from: <https://guidelines.diabetes.ca/cpg/chapter13>
 66. About the National Health and Nutrition Examination Survey [Internet]. 2017 [cited 2021 Aug 29]. Available from: https://www.cdc.gov/nchs/nhanes/about_nhanes.htm
 67. CDC. Body Mass Index (BMI) [Internet]. 2021 [cited 2021 Sep 2]. Available from: <https://www.cdc.gov/healthyweight/assessing/bmi/index.html>

68. Anthropometry Procedures Manual [Internet]. CDC Centers for Disease Control and Prevention. 2017 [cited 2021 Sep 2]. p. 1-1-5–14. Available from: https://wwwn.cdc.gov/nchs/data/nhanes/2017-2018/manuals/2017_Anthropometry_Procedures_Manual.pdf
69. How the Census Bureau Measures Poverty [Internet]. United States Census Bureau. 2020 [cited 2021 Aug 28]. Available from: <https://www.census.gov/topics/income-poverty/poverty/guidance/poverty-measures.html#:~:text=The total family income divided by the poverty,Income – Threshold %3D %2432%2C000 - %2431%2C275%3D %24725>
70. Schernthaner G, Matthews DR, Charbonnel B, Hanefeld M, Brunetti P. Efficacy and safety of pioglitazone versus metformin in patients with type 2 diabetes mellitus: A double-blind, randomized trial. *J Clin Endocrinol Metab*. 2004;89(12):6068–76.
71. Edelman S V. Optimizing diabetes treatment using an amylin analogue. *Diabetes Educ*. 2008;34(1):4–10.
72. Fiorentino TV, Sesti G. Lessons learned from cardiovascular outcome clinical trials with dipeptidyl peptidase 4 (DPP-4) inhibitors. *Endocrine*. 2016;53(2):373–80.
73. Smilowitz NR, Donnino R, Schwartzbard A. Glucagon-like peptide-1 receptor agonists for diabetes mellitus: A role in cardiovascular disease. *Circulation*. 2014;129(22):2305–12.
74. Hermann LS, Scherstén B, Bitzén P-O, Kjellström T, Lindgärde F, Melander A. Therapeutic Comparison of Metformin and Sulfonylurea, Alone and in Various Combinations: A double-blind controlled study. *Diabetes Care* [Internet]. 1994 Oct 1;17(10):1100 LP – 1109. Available from:

<http://care.diabetesjournals.org/content/17/10/1100.abstract>

75. Rosenstock J, Hassman D, Maddar R, Farrel J, Khutoryansky N. Repaglinide versus nateglinide monotherapy: a randomized, multicenter study. *Diabetes Care*. 2004;27(6):1265+.
76. Klein S, Sheard NF, Pi-Sunyer X, Daly A, Wylie-Rosett J, Kulkarni K, et al. Weight management through lifestyle modification for the prevention and management of type 2 diabetes: Rationale and strategies. A statement of the American Diabetes Association, the North American Association for the Study of Obesity, and the American So. *Am J Clin Nutr*. 2004;80(2):257–63.
77. Lee JJ, Beretvas SN, Freeland-Graves JH. Abdominal adiposity distribution in diabetic/prediabetic and nondiabetic populations: A meta-analysis. *J Obes*. 2014;2014.
78. Sathiyapriya V, Selvaraj N, Nandeesh H, Bobby Z, Agrawal A, Sridhar MG, et al. Increased glycation of hemoglobin and plasma proteins in normotensive, non-diabetic obese Indian subjects: Putative role of lipid peroxides. *Clin Chem Lab Med*. 2007;45(8):996–9.
79. Hodish I. Insulin therapy, weight gain and prognosis. *Diabetes, Obes Metab*. 2018;20(9):2085–92.
80. Costella R, Nicolas S, Shivkumar A. Sulfonyleureas. StatPearls Publishing; 2021.
81. Mc Auley MT. Effects of obesity on cholesterol metabolism and its implications for healthy ageing. *Nutr Res Rev*. 2020;33(1):121–33.
82. Saleem Z, Saeed H, Khan ZA, Khan MIH, Hashmi FK, Islam M, et al. Association of

- hypertension and dyslipidaemia with increasing obesity in patients with type 2 diabetes mellitus. *Brazilian J Pharm Sci.* 2019;55:1–9.
83. Kovanen PT, Nikkila EA, Miettinen TA. Regulation of cholesterol synthesis and storage in fat cells. *J Lipid Res.* 1975;16(3):211–23.
84. Krause BR, Hartman AD. Adipose tissue and cholesterol metabolism. *J Lipid Res* [Internet]. 1984;25(2):97–110. Available from: [http://dx.doi.org/10.1016/S0022-2275\(20\)37830-5](http://dx.doi.org/10.1016/S0022-2275(20)37830-5)
85. Wulffelé MG, Kooy A, De Zeeuw D, Stehouwer CDA, Gansevoort RT. The effect of metformin on blood pressure, plasma cholesterol and triglycerides in type 2 diabetes mellitus: A systematic review. *J Intern Med.* 2004;256(1):1–14.
86. Sharma AM. Obesity and cardiovascular risk. *Growth Horm IGF Res.* 2003;13(SUPPL. A):10–7.
87. Mokdad AH, Ford ES, Bowman BA, Dietz WH, Vinicor F, Bales VS, et al. Prevalence of obesity, diabetes, and obesity-related health risk factors, 2001. *J Am Med Assoc.* 2003;289(1):76–9.
88. Andreyeva T, Michaud PC, van Soest A. Obesity and health in Europeans aged 50 years and older. *Public Health* [Internet]. 2007;121(7):497–509. Available from: <http://dx.doi.org/10.1016/j.puhe.2006.11.016>
89. Bauman AE, Grunseit AC, Rangul V, Heitmann BL. Physical activity, obesity and mortality: Does pattern of physical activity have stronger epidemiological associations? *BMC Public Health.* 2017;17(1):1–12.

90. Goodpaster BH, DeLany JP, Otto AD, Kuller L, Vockley J, South-Paul JE, et al. Effects of diet and physical activity interventions on weight loss and cardiometabolic risk factors in severely obese adults: A randomized trial. *JAMA - J Am Med Assoc.* 2010;304(16):1795–802.
91. Canning KL, Brown RE, Jamnik VK, Kuk JL. Relationship between obesity and obesity-related morbidities weakens with aging. *Journals Gerontol - Ser A Biol Sci Med Sci.* 2014;69(1):87–92.
92. Woolley C, Thompson C, Hakendorf P, Horwood C. The Effect of Age upon the Interrelationship of BMI and Inpatient Health Outcomes. *J Nutr Heal Aging.* 2019;23(6):558–63.
93. Woo J, Ho SC, Yu ALM, Sham A. Is waist circumference a useful measure in predicting health outcomes in the elderly? *Int J Obes.* 2002;26(10):1349–55.
94. Blair SN, Church TS. Is Physical Activity the Common Denominator ? *JAMA [Internet].* 2004;292(10):12321234. Available from: <http://jama.jamanetwork.com/article.aspx?articleid=199376>
95. Lee DC, Sui X, Blair SN. Does physical activity ameliorate the health hazards of obesity? *Br J Sports Med.* 2009;43(1):49–51.
96. Colberg SR. Key points from the updated guidelines on exercise and diabetes. *Front Endocrinol (Lausanne).* 2017;8(FEB):1–7.
97. Zhao G, Ford ES, Li C, Balluz LS. Physical activity in U.S. older adults with diabetes mellitus: Prevalence and correlates of meeting physical activity recommendations. *J Am*

- Geriatr Soc. 2011;59(1):132–7.
98. Micha R, Shulkin ML, Peñalvo JL, Khatibzadeh S, Singh GM, Rao M, et al. Etiologic effects and optimal intakes of foods and nutrients for risk of cardiovascular diseases and diabetes: Systematic reviews and meta-analyses from the nutrition and chronic diseases expert group (NutriCoDE). PLoS One. 2017;12(4):1–25.
 99. Noakes M, Clifton P, McMurchie T. The role of diet in cardiovascular health. A review of the evidence [corrected] [published erratum appears in AUST J NUTR DIET 1999 Dec;56(4):186]. Aust J Nutr Diet [Internet]. 1999;56(3):S3-22. Available from: <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=2003021182&site=ehost-live>
 100. Diabetes and Meal Planning [Internet]. CDC Centers for Disease Control and Prevention. 2021. Available from: <https://www.cdc.gov/diabetes/managing/eat-well/meal-plan-method.html>
 101. Diabetes diet: Create your healthy-eating plan [Internet]. Mayo Clinic. 2021. Available from: <https://www.mayoclinic.org/diseases-conditions/diabetes/in-depth/diabetes-diet/art-20044295>
 102. Mohammed MA, Sharew NT. Adherence to dietary recommendation and associated factors among diabetic patients in ethiopian teaching hospitals. Pan Afr Med J. 2019;33:1–11.
 103. Wild S, Roglic G, Green A, Sicree R, King H. Global Prevalence of Diabetes: Estimates for the year 2000 and projections for 2030. Diabetes Care. 2004;27(5):1047–53.

104. Chi C, Snaith J, Gunton JE. Diabetes Medications and Cardiovascular Outcomes in Type 2 Diabetes. *Hear Lung Circ* [Internet]. 2017;26(11):1133–41. Available from: <http://dx.doi.org/10.1016/j.hlc.2017.02.030>
105. Williams GC, Patrick H, Niemiec CP, Williams LK, Divine G, Lafata JE, et al. Reducing the health risks of diabetes: How self-determination theory may help improve medication adherence and quality of life. *Diabetes Educ*. 2009;35(3):484–92.
106. Capoccia K, Odegard PS, Letassy N. Medication Adherence With Diabetes Medication: A Systematic Review of the Literature. *Diabetes Educ*. 2016;42(1):34–71.