

Stable carbon isotope signatures of background tropospheric chloromethane and CFC113

ALEXANDRA E. THOMPSON^{1,*}, REBECCA S. ANDERSON¹, JOCHEN RUDOLPH¹ & LIN HUANG²

Key words: CFC113, CH₃Cl, GC-IRMS, halocarbons, stable carbon isotope ratios, tropospheric chloromethane

Abstract. Samples of background air were collected in the lower troposphere of the Northern (high Arctic, northern Ontario, Vancouver and Houston) and Southern (Baring Head, New Zealand) Hemispheres over the period July 1999 until March 2001. These samples were analysed for the stable carbon isotope ratios of 1,1,1-trichlorotrifluoroethane (CFC113) and CH₃Cl using a gas chromatography-continuous flow on-line combustion isotope ratio mass spectrometry combination. For CH₃Cl the global average of the stable carbon isotope ratio is $-36.2 \pm 0.3\text{‰}$ (error of mean). The average is based on 78 data points, standard deviation for all measurements is 2.3‰, and the 90% confidence interval is -35.8 to -36.6‰ . However, the number of data points from the Southern Hemisphere is rather limited and thus this observation is not necessarily representative for the entire Southern Hemisphere. A simple isotopic budget of CH₃Cl shows the most important parameters needing to be defined are the kinetic isotope effect of CH₃Cl destruction by OH radicals and the source composition of CH₃Cl emitted by the oceans and biomass burning of C-4 plants. Present budgets of atmospheric CH₃Cl show a significant deficit in the source strength. We estimate that the average stable carbon isotope ratio of the additional CH₃Cl emissions required to balance the budget is $-41.9 \pm 7.8\text{‰}$. The average CFC113 isotopic composition based on 38 measurements is $-23.3 \pm 1.6\text{‰}$ (error of mean), $\sigma = 9.6\text{‰}$ with no significant difference between the hemispheres.

Introduction

Chloromethane (CH₃Cl) with an average atmospheric concentration in the range of 500–600 ppt represents one of the largest reservoirs of gaseous chlorine in the atmosphere. It contributes 43% of the total reactive Cl in the troposphere and is increasingly important as a source for Cl atoms in the stratosphere (Khalil et al. 1999, Khalil & Rasmussen 1999). It comprises the

only significant source of natural Cl atoms to the stratosphere, contributing up to 25% (Keene et al. 1999) of stratospheric Cl. Since the emission rates of trace gases can be influenced by various factors, a quantitative understanding of the atmospheric CH₃Cl budget is essential in order to allow predictions of potential future developments of CH₃Cl emissions, e.g. as a consequence of climate change.

However, our current understanding of the global budget of tropospheric CH₃Cl is completely unsatisfactory. The quantified sources can only account for approximately half of the emissions required to balance the known sinks. The major sink of CH₃Cl in the troposphere is loss by reaction with OH-radicals (3.37 Tg CH₃Cl/yr). Minor sinks include loss to the stratosphere and uptake by soils (0.28 and 0.25 Tg CH₃Cl/yr respectively). Current estimates of emissions only total 2.03 Tg CH₃Cl/yr (Keene et al. 1999; Rhew et al. 2000), which is approximately 50% of the atmospheric loss rate.

The most important known CH₃Cl source is biomass burning, contributing 0.90 Tg CH₃Cl annually. Oceanic emissions are estimated to account for 0.65 Tg CH₃Cl/yr, salt marsh emissions 0.17 Tg CH₃Cl/yr (Rhew et al. 2000), and fungi, coal combustion, incineration and industry emit a total of 0.31 Tg CH₃Cl/yr (Keene et al. 1999).

Clearly, either the emissions from the identified sources are underestimated, the sinks are overestimated, or there are other unknown sources of CH₃Cl. Since the dominant atmospheric loss process of CH₃Cl, reaction in the troposphere with OH radicals, is well understood, an overestimate of the loss rates is an unlikely explanation for the very substantial imbalance in the CH₃Cl budget.

Stable carbon isotope ratio measurements have been used very successfully to constrain the atmospheric budgets of a number of important trace gases such as carbon dioxide, carbon monoxide, and methane (e.g. Kaplan 1975; Lowe et al. 1994; Mook et al. 1983). With the recent development of techniques to measure the stable carbon isotope ratio of volatile organic compounds in the atmosphere (Rudolph et al. 1997) the use of isotope composition measurements to improve our insight into the atmospheric cycle of CH₃Cl became feasible. However, there is still very little published data available on the stable carbon isotope ratios of ambient CH₃Cl. Knowledge of a representative average tropospheric distribution of the stable carbon isotope ratio of CH₃Cl is an essential prerequisite for any attempt to use stable carbon isotope studies to improve our present understanding of the global CH₃Cl budget. Currently there are only two published papers on ambient CH₃Cl isotopic composition (Tsunogai et al. 1999; Rudolph et al. 1997). Rudolph et al. presented one measurement from a background station in New Zealand, Tsunogai et al. made measurements in urban and coastal areas of Japan and

over the Pacific Ocean between 5° N and 35° N latitude. These measurements gave stable carbon isotope ratios of CH₃Cl in the range of -30 to -45‰ ($\delta^{13}\text{C}$ relative to Pee Dee Belemnite). Obviously, the limited geographic coverage of the available ambient measurements and the substantial spread of the data will severely limit the reliability of any budget constraints based on the presently existing ambient data. In this paper we present the results from seven campaigns comprising a total of 90 samples that allows significant reduction of the uncertainty of the average global stable carbon isotope ratio of CH₃Cl.

We also measured the isotopic composition of 1,1,1-trichlorotrifluoroethane (CFC113). CFC113 is a manufactured compound that was used in refrigeration, as a solvent, and a degreasing agent before its ban by the Montreal Protocol. In 1992 it was present in the atmosphere at a concentration of 82 ppt and had an atmospheric burden of 2.6 Tg (IPCC 1995). Present emissions are negligible and the dominant tropospheric loss process is transport to the stratosphere and subsequent photolysis. There is nothing published on the isotopic composition of CFC113 to date.

Experimental procedure

Whole air samples were collected in the field and transferred for analysis to the laboratory. Following a cryogenic enrichment procedure, the stable carbon isotope ratios of CH₃Cl and CFC113 were determined by isotope ratio mass spectrometry in combination with a gas chromatographic separation and continuous flow on-line combustion. The procedure was very similar to a method described by Rudolph et al. (1997), but a few modifications were made and therefore a brief description of the technique is presented here.

Ambient Sampling

Sampling locations, characterisation of sampled air masses, number of samples, sampling periods, and sampling frequency are tabulated in Table 1. The ambient air samples were collected in evacuated electropolished 3L stainless steel canisters at a pressure of approximately 2 atm above ambient. For background samples, as well as the aircraft samples, the canisters were pressurised using metal bellows pumps in order to minimise the risk of sample contamination. Tests showed that metal bellows pumps did not cause detectable levels of contamination for any of the compounds studied here. For urban or semi-urban samples Teflon diaphragm pumps were used. For these samples of somewhat higher trace gas levels the use of Teflon diaphragm pumps caused no detectable change in the trace gas concentration. The advantage

of Teflon diaphragm pumps is their lower power consumption and thus the possibility to use batteries as power supply. This increased the flexibility in selection of sampling locations.

Pre-concentration

A two step process was used to enrich the analytes from between 1 and 20 L of air. Prior to the enrichment steps CO₂ was removed by passing the sample air through a 30 cm × 1/2" stainless steel tube filled with Ascarite II. In order to maintain the CO₂ removal efficiency at a high level (>99.9%), the Ascarite trap was conditioned prior to each measurement by passing humidified He at a flow rate of 30 cm³ min⁻¹ for 30 minutes through the trap. After every second analysis the trap was replaced by a new trap containing fresh Ascarite. Prior to use all traps were conditioned for at least 8 hours in a He flow at 80 °C to eliminate contamination. Testing of this procedure (including the introduction of a CH₃Cl standard before and after the trap) showed that there were no losses or fractionations for CH₃Cl and that CO₂ removal is highly efficient (>99.9%) for sample volumes of 30 L or less. It must be noted that no tests of the analytical method specific for CFC113 were performed. Thus all stable carbon isotope ratios presented for CFC113 are based on the assumption that the tests conducted for CH₃Cl were also valid for CFC113.

After the CO₂ trap, the sample flowed through a two-stage water trap (15 cm × 1" stainless steel and then 50 cm × 3/8" stainless steel, both held at -30 °C during pre-concentration) before passing through the first cryogenic trap. In this stainless steel trap of 30 cm length and 3/8" diameter packed with 80/100 mesh glass beads, the volatile organic compounds in the sample were cryogenically trapped at liquid Ar temperature (-186 °C). Sample flow rate through this trap was 100 ml/min. Consequently, for a 15 L sample this process took 2.5 hours. However, the sample enrichment procedure was usually performed during the chromatographic separation of the previous sample to reduce the overall analysis time. The trap was then heated to 160 °C and the halocarbons transferred in a flow of 5-12 ml/min of He to a second trap (8 cm × 1/16" stainless steel) held at -196 °C with liquid N₂ for cryogenic focussing. For sample injection this trap was flash heated and the pre-concentrated halocarbons were transferred to the separation columns by the carrier gas.

Separation and analysis

The samples were separated on a Varian Series 3600 gas chromatograph using a combination of a 60 m (0.32 mm ID, 5 μm film) HP1 column and a 60 m (0.32 mmID) Chromatographic Specialties Poraplot Q (PPQ) column. Flow

Table 1. Overview of sampling campaigns

Campaign	Location	Number of Samples	Number of Samples of CH3CI	Number of Samples of CFCl13	Characterisation of sample air	Sampling Period	Frequency
Alert, seasonal cycles	Alert, Nunavut, Canada (82.5°N, 62.3°W)	42	31		Background	Jul 1999–Mar 2001	One sample every fortnight
Fraserdale	Fraserdale, Ontario, Canada (49.9°N, 81.6°W)	12	0		Background, continental	Apr 2000–Feb 2001	One sample every three weeks
Vancouver	Fraser Valley, British Columbia, Canada (48.3°N, 123.7°W)	1	0		Rural	5 May 2000	
Houston	Houston, Texas, United States (29.8°N, 95.3°W)	16	0		Background into urban plumes, aircraft campaign	29, 31 Aug 2000	3 flights
New Zealand	Baring Head, New Zealand (41.4°S, 174.9°E)	16	5		Background	Jul–Sep 1999	Approx. one sample every three weeks
New Zealand	Auckland, New Zealand (36.8°S, 175.7°E)	2	0		Suburban, close to motorway	3, 14 March 2000	
Polar Sunrise 1999	Alert, Nunavut, Canada (82.5°N, 62.3°W)	0	2		Background	3, 20 May 1999	

rates were 1.5 ml/min, the carrier gas was He. The sample was pre-separated on the HP1 column and the fraction of the column effluent containing CH₃Cl was transferred to the PPQ column for further separation. CFC113 was separated on the HP1 column only. The temperature was held at -30 °C for 30 mins, then a 3 °C/min ramp to 180 °C was applied. About 0.5 ml/min of the column effluent was transferred via a deactivated fused silica capillary to an Ion Trap mass spectrometer for peak identification and verification of peak purity. The remainder of the carrier gas flow entered a commercial (Finnigan MAT) combustion interface similar to that described by Matthews and Hayes (1978) to convert the analyte into CO₂, and about 0.3 ml/min of the flow was transferred via an open split to a Finnegan Mat 252 Isotope Ratio Mass Spectrometer (IRMS). The isotopic composition of the CO₂ from each individual compound was measured against a CO₂ reference gas (with $\delta^{13}\text{C}_{\text{VPDB}} = -35.4\text{‰}$) from the reference open split. This on-line reference was calibrated to the primary scale Vienna Pee Dee Belemnite (VPDB) scale on a daily basis using two lab carbonate standards with $\delta^{13}\text{C}_{\text{VPDB}}$ of -2.57‰ and -45.78‰ . All delta values in this paper are reported relative to this scale, and presented errors, unless otherwise stated, are the standard error of the mean.

Results

Chloromethane

Concentration

CH₃Cl concentrations were not measured separately in any of the campaigns. However, for the Alert data set the CH₃Cl concentrations have been determined from the 44 mass trace peak area of CH₃Cl relative to the 44 mass trace peak area of propane, for which the concentration was measured independently using GC-FID methods. CH₃Cl and propane underwent separation on the same columns and elute within 50s of each other. We estimate the error in the CH₃Cl concentration is 10%, mainly due to the uncertainty of the propane concentration, and the possibility of minor changes in the column flow rate between elution of the CH₃Cl peak and the propane peak, which effect the fraction of analyte reaching the IRMS. The CH₃Cl concentrations are shown in Figure 1 along with the 7-year monthly average at Alert during the period 1991–1996 (Gautrois et al. 2002). Yokouchi et al. (1998) also measured CH₃Cl at Alert and observed similar concentrations, although the magnitude of the seasonal variation was slightly less than that of Gautrois et al. According to both of these studies, CH₃Cl at Alert has an average concentration of 500 ppt and shows a seasonal variation of approximately 150–230 ppt with a maximum concentration in winter. Within the uncer-




Figure 1. Concentration of CH_3Cl at Alert. Solid squares represent concentrations from this work; open circles represent monthly averages from observations during 1991 to 1996 (Gautrois et al. in prep.). The error bars represent the measurement error, and the error of the mean, respectively.

tainty of measurements and the expected interannual variability, our CH_3Cl concentrations are compatible with these other studies. Unfortunately it was not possible to calculate the concentrations for other campaigns using this method because the chromatography differed and associated errors became too large.

Isotopic composition

Table 2 summarises the CH_3Cl stable carbon isotopic composition data. The average isotopic composition of background tropospheric CH_3Cl in the Northern Hemisphere from our observations is $-36.1 \pm 0.3\%$, $N = 68$.

The stable carbon isotopic composition and the concentrations for CH_3Cl at Alert are plotted against each other in Figure 2. The plot shows no correlation between isotopic composition and concentration ($R^2 = 0.001$), however there are three clearly outlying data points with concentrations greater than 1.4 ppb . These data, likely to be contaminated or influenced by unusually high local or regional emissions, have been removed from the data set.

The seasonal averages are shown in Table 3. The average summer (June until August) composition is $-36.5 \pm 0.7\%$, $N = 9$, the average for winter (December until February) is $-35.8 \pm 0.4\%$, $N = 11$, and the annual average is $-36.4 \pm 0.3\%$, $N = 39$ at Alert. Within their statistical uncertainty of less than 1% these values are the same. At Fraserdale the average winter composition, $-35.0 \pm 0.7\%$, $N = 3$, and the annual average is $-36.0 \pm 0.6\%$, $N =$

Table 2. Average isotopic composition of CH₃Cl (all values in ‰^b)

Campaign	Alert ^b	Fraserdale	Vancouver	Houston	New Zealand ^c	New Zealand Background ^d	New Zealand Urban	Northern Hemispheric Average	Global Average
Average δ ¹³ C	-36.40	-35.99	-38.88	-35.17	-42.13	-37.23	-40.64	-36.08	-36.22
Error of mean	0.26	0.64	0.5 ^e	0.83	1.85	0.82	8.90	0.27	0.26
Standard Deviation	1.63	2.12	-	3.20	7.18	2.47	8.90	2.22	2.27
Median	-36.62	-35.75	-	-35.34	-39.00	-37.79	-40.64	-36.37	-36.61
25th Percentile	-37.67	-37.01	-	-37.42	-47.04	-38.56	-43.79	-37.56	-37.69
75th Percentile	-35.53	-34.57	-	-34.00	-37.56	-36.75	-37.50	-34.90	-35.01
90% Confidence Interval	±0.43	±1.01	-	±1.32	±2.95	±1.28	±10.35	±0.44	±0.42
Number of samples	39	12	1	16	16	10	2	68	78

^aRelative to VPDB.

^bExcluding contaminated samples.

^cIncluding all samples taken in New Zealand Background campaign.

^dExcluding samples possibly directly impacted by recent emissions.

^eError of measurement.




Figure 2. Concentrations and isotopic composition of CH₃Cl at Alert.

12, are also the same as the averages of our measurements at Alert. Only the stable carbon isotope ratio for summer, with $-38.5 \pm 1.4\%$, $N = 3$, is slightly lower than our observations at Alert. However this summer average is based on a very small number of measurements. Our other background tropospheric CH₃Cl stable carbon isotope measurements in the Northern Hemisphere fall into a very similar range (Table 2). The average is $-36.1 \pm 0.3\%$, $N = 68$. The data median is -36.4% , and the confidence interval at the 90% level is -35.6% to -36.5% . The 25th and 75th percentiles are -37.6% and -34.9% respectively. This is within the range of results previously found by Tsunogai et al. (1999) of -44 to -30% for marine and coastal CH₃Cl, but our measurements allow the determination of a considerably better defined representative average. There is no statistically significant difference between background samples from the different locations at different latitudes in the Northern Hemisphere.

The stable isotope ratios of CH₃Cl in background samples from the Southern Hemisphere show a less uniform picture (Table 2). The available data are limited to a relatively small region, New Zealand, and show a larger variability than the Northern Hemispheric measurements. The average of all our Southern Hemispheric measurements is $-42.1 \pm 1.9\%$, $N = 16$, noticeably lighter than our Northern Hemispheric average, however the substantial difference between the median and the mean (Table 2) indicates a somewhat skewed distribution. The frequency distributions for the Northern Hemispheric and New Zealand stable carbon isotope ratio measurements are compared in Figure 3. Data from the Northern Hemisphere present an approximately symmetrical frequency distribution, however the New Zealand

Table 3. Seasonal averages of isotopic composition of CH₃Cl in two Northern Hemispheric background sites (all values in ‰^a)

Season	Summer (Jun–Aug)	Autumn (Sep–Nov)	Winter (Dec–Feb)	Spring (Mar–May)	Seasonal Variation Winter– Summer
Campaign	Alert	Alert	Alert	Alert	Alert
Average $\delta^{13}\text{C}$	-36.53	-36.28	-35.83	-37.35	0.69
Error of mean ^b	0.69	0.56	0.37	0.50	0.78
Standard deviation	1.94	1.86	1.17	1.22	
Number of samples	9	12	11	7	
Campaign	Fraserdale	Fraserdale	Fraserdale	Fraserdale	Fraserdale
Average $\delta^{13}\text{C}$	-38.50	-34.74	-35.02	-36.19	3.47
Error of mean ^b	1.44	1.03	0.71	1.20	1.61
Standard deviation	2.04	1.78	1.00	1.20	
Number of samples	3	4	3	2	

^arelative to VPDB

^bfor seasonal variation this value is the error of the difference.

samples do not. Since the atmospheric lifetime of CH₃Cl (1.3 years) is substantially greater than the hemispheric mixing time (3 months) the hemispheric background troposphere should be well mixed with respect to CH₃Cl and therefore the distribution of delta values should be narrow. The wide range of observed values indicates that some of the samples from New Zealand must have been impacted by relatively recent CH₃Cl emissions, which are depleted in ¹³C. Biomass burning in the tropics and subtropics of the Southern Hemisphere is a very important source of CH₃Cl (Andreae 1991). Rudolph et al. (1997) and von Czapiewski et al. (2001) measured the isotopic composition of CH₃Cl of biomass burning. Both studies show that CH₃Cl emitted from biomass burning is relatively light, $\delta^{13}\text{C}$ values range from -38‰ to -68‰. Other known emissions of CH₃Cl for which the isotopic composition has been measured also have low $\delta^{13}\text{C}$ values (Table 4). Obviously, even a very moderate impact of emissions of very light CH₃Cl into the studied air mass might decrease its ambient stable carbon isotope ratio by several per mil. Our Southern Hemispheric data do not allow any conclusions on the representativeness of the unusually low values. If we exclude these data from the average, we obtain a value of $-37.2 \pm 0.8\text{‰}$, $N = 10$. This value, within its statistical uncertainty, is the same as the Northern Hemispheric average. Our data thus indicate that CH₃Cl in the Southern Hemisphere has




Figure 3. Frequency distributions of the isotopic composition of CH_3Cl in the (a) Southern Hemisphere and (b) Northern Hemisphere data sets.

a stable carbon isotope ratio very similar to that of the Northern Hemisphere, although there are a number of observations that indicate depletion in ^{13}C relative to the Northern Hemisphere. The suburban New Zealand samples had CH_3Cl isotopic compositions of $-46.9 \pm 0.5\%$ and $-34.4 \pm 0.5\%$.

CFC113

The measured CFC113 isotopic compositions are shown in Table 5. The global average stable carbon isotopic composition of ambient CFC113 is $23.3 \pm 1.6\%$, $N = 38$. The Alert campaign was analysed with a slightly different chromatographic separation, which caused occasional partial peak merging with an unknown species. This interference did not occur in either the Polar Sunrise 1999, or the New Zealand Background measurements. This analytical problem is responsible for the high variability in the Alert data set.

Table 4. Known Sources of Tropospheric CH₃Cl

Sources	Source Strength ^a (TgCH ₃ Cl/yr)	Source Composition (%)	Error in Composition (%)
Biomass burning	0.90	-47 ^b	12 ^b
Fungi	0.15	-43 ^c	2 ^c
Salt Marshes	0.17 ^d	-62 ^e	3 ^e
Industry	0.01	-41.9 and -60 ^f	
Ocean	0.65	?	
Coal combustion	0.11	?	
Incineration	0.04	?	
TOTAL	2.03		

^aKhalil et al. 1999a.

^bWeighted average of C3 and C4 plants, see text for details.

^cHarper et al. 2001, not including higher plants.

^dRhew et al. 2000.

^eBill et al. 2002.

^fComposition of manufactured CH₃Cl (Rudolph et al. 1997 and Tsunogai et al. 1999).

Table 5. Average isotopic composition of CFC113 (all values in ‰^a)

Campaign	Polar Sunrise 1999	Alert Zealand	New Average of Background	Global Sample Ambient CFC113	Commercial
Average δ ¹³ C	-22.70	-23.53	-22.24	-23.32	-31.3
Error of mean	1.27	1.95	0.66	1.58	0.5 ^b
Standard Deviation	1.27	10.69	1.32	9.64	-
Median	-22.70	-24.74	-23.10	-23.10	-
25th Percentile	-23.15	-29.08	-23.10	-28.58	-
75th Percentile	-22.25	-21.28	-21.80	-21.55	-
90% Confidence interval	±1.48	±3.15	±0.97	±2.57	-
Number of samples	2	31	5	38	1

^aRelative to VPDB.

^bEstimated measurement error.

The Northern Hemispheric average stable isotopic composition is $-23.5 \pm 1.8\text{‰}$ N=33, and the Southern Hemispheric average is $-22.2 \pm 0.7\text{‰}$, $N = 5$. There is no significant difference in the isotopic composition of ambient CFC113 between the hemispheres. For comparison, a commercial sample (Sigma Aldrich Lot # B100453TU, produced in December 1999 by Aldrich) of CFC113 was analysed and its stable carbon isotope ratio was found to be $-31.3 \pm 0.5\text{‰}$.

Discussion

Chloromethane

Seasonal variation in the concentration of CH_3Cl has been observed in the high and mid latitudes and the tropics of both hemispheres (although at different magnitudes) and is the result of the seasonal variation of the OH radical concentrations (Khalil & Rasmussen 1999b). The kinetic isotope effect (KIE) of CH_3Cl and OH is not known. To date, all measured KIEs of hydrocarbons reacting with OH have been positive (Rudolph et al. 2000). Assuming that the KIE of CH_3Cl and OH is positive, as atmospheric CH_3Cl is photochemically aged its isotopic composition becomes heavier, i.e. enriched in ^{13}C . Thus, we expect that it would be heavier in summer than during the winter. However, the seasonal variation seen in the Fraserdale data set shows the winter isotopic composition to be slightly heavier than the summer composition. This is an indication that CH_3Cl at Fraserdale is impacted by lighter sources during the summer than during the winter. Thus the seasonal variation at Fraserdale can be explained by a regional impact of biomass burning. A seasonal variation of 3.5‰ would only require a 15% additional contribution from biomass burning during the summer. Summertime biomass burning is known to occur at mid and boreal latitudes, and there is substantial burning in Canada. Further away from the sources, for example at Alert where the burning forests are at least 1000 km or more away, the effect will be diluted.

It is possible to construct a simple global budget for CH_3Cl using the isotopic compositions and associated errors to constrain the contribution of different sources. However, substantial difficulties lie with firstly the parameters which are completely unknown, and secondly that the uncertainties of the 'known' parameters are not understood. Most notably, errors in the magnitude of the sources and sinks of tropospheric CH_3Cl are, for the most part, undefined. Although accuracy in the magnitude of sources and sinks is critical to any budget, for the purposes of this paper we have used the most likely magnitudes and neglected their associated errors in the budget calcula-

tions. This is a substantial weakness in the discussion of the isotopic budget of CH_3Cl , however since so little is understood about the sources of atmospheric CH_3Cl , stable carbon isotope arguments can still be useful. There are also many gaps in our understanding of the effect of specific sources and sinks on the isotopic composition of atmospheric CH_3Cl . We use 'best guess' estimations of the effects and errors not explicitly stated in the literature.

An overview of identified CH_3Cl sources and their isotopic compositions is shown in Table 4. As previously mentioned, only about half of the emissions required to balance the atmospheric CH_3Cl budget have been identified. The most important known chloromethane source is biomass burning which emits 0.90 Tg $\text{CH}_3\text{Cl}/\text{yr}$ (48% of known sources). As mentioned above, the two existing studies (Rudolph et al. 1997; von Czapiewski 2001) indicate that chloromethane emitted from biomass burning has stable carbon isotope ratios from -38‰ to -68‰ , with an average of -52‰ . However, both published studies used material from plants with a C-3 metabolism as fuel. Plants with a C-4 metabolism are enriched in ^{13}C relative to C-3 plants (parent material of -13‰ and -27‰ respectively) and are thought to be responsible for about a third of CH_3Cl emissions from biomass burning (Lobert et al. 1999). Isotopic fractionation during combustion of C-3 plants is $-25 \pm 12\text{‰}$. We assume that the fractionation relative to the parent fuel is the same for the combustion of material derived from C-3 and C-4 plants, that is $-25 \pm 12\text{‰}$. The resulting average stable carbon isotope ratio of CH_3Cl from biomass burning is $-47 \pm 12\text{‰}$.

Current estimates place oceanic emissions of CH_3Cl to be the second largest known source at 0.65 Tg $\text{CH}_3\text{Cl}/\text{yr}$, about 70% of the size of biomass burning emissions. Unfortunately, nothing is known about the stable carbon isotope ratio of CH_3Cl produced and emitted by oceans. The composition of oceanic CH_3Cl is unknown, and although fixed carbon in seawater is known to have a delta value of -22‰ , the isotopic composition of oceanic emissions of CH_3Cl is not necessarily representative of the source material.

Other known sources include fungi (0.15 Tg $\text{CH}_3\text{Cl}/\text{yr}$), salt marshes (0.17 Tg $\text{CH}_3\text{Cl}/\text{yr}$), and coal combustion, incineration, and industry (total of 0.16 Tg $\text{CH}_3\text{Cl}/\text{yr}$). CH_3Cl produced by one species of fungi was measured to be -43.3‰ (Harper et al. 2001). The isotopic composition of emissions of CH_3Cl from salt marshes ranges from -45‰ to -71‰ , with a daily weighted mean of $-62 \pm 3\text{‰}$ (Bill et al. 2002). Industrial CH_3Cl has been measured to be -60‰ and -41.9‰ (Tsunogai et al. 1999; Rudolph et al. 1997). The isotopic composition of CH_3Cl emissions from coal combustion and incineration are not known. Overall the isotopic composition of these minor sources is extremely uncertain. However, they contribute only about 10% to the total CH_3Cl budget.

Table 6. Known Sinks of Tropospheric CH₃Cl

Sinks	Sink Magnitude ^a (TgCH ₃ Cl/yr)	Fractionation Factor, ϵ (‰)	Error in ϵ (‰)
OH Destruction	3.37	5 ^b	5 ^b
Loss to Stratosphere	0.28	5 ^b	5 ^b
Soils	0.25	46 ^c	20 ^b
TOTAL	3.91		

^aKhalil et al. 1999a.

^bEstimate, see text for details.

^cMüller et al. 2001.

The known CH₃Cl sink mechanisms and their associated fractionation factors as well as uncertainties are summarised in Table 6. The total loss rate is 3.91 Tg CH₃Cl/yr. The dominant loss process of atmospheric CH₃Cl by far is via reaction with OH-radicals (3.37 Tg CH₃Cl/yr). Currently the kinetic isotope effect (KIE) for reaction of CH₃Cl with the OH radical is unknown.

Another loss of atmospheric chloromethane is the destruction by methylotrophic bacteria in soils. It is a minor sink of 0.25 Tg CH₃Cl/yr (6.5% of total loss), however it potentially has a significant fractionation effect. Müller et al. 2001 measured the fractionation of CH₃Cl to be 46‰ for three strains of (whole cell) bacteria at elevated methyl halide concentrations. Although it is not clear how representative this study is to the effects on ambient atmospheric CH₃Cl, this fractionation factor has been included in the budget with an estimated error of 20‰.

And finally, 0.28 Tg CH₃Cl/yr of tropospheric CH₃Cl is known to be lost to the stratosphere where it is destroyed by photolysis. Since this CH₃Cl sink is via turbulent mixing, the transport process itself will not result in an isotopic fractionation. It cannot be excluded that stratospheric loss, which is via photolysis and reaction with OH-radicals, will be associated with an isotope fractionation. However, stratospheric loss is less than 10% of tropospheric removal and it is therefore extremely unlikely that this loss has a major impact on the isotopic composition of tropospheric CH₃Cl. We use the same fractionation and uncertainty for the stratospheric and tropospheric loss processes.

The global average tropospheric CH₃Cl composition is equal to the weighted average of all emissions plus the weighted average of removal mechanisms (equation 1).

$$\delta_{obs} = \frac{\sum_{all-sources} \delta_i * S_i}{\sum_{all-sources} S_i} + \frac{\sum_{all-losses} \epsilon_j * L_j}{\sum_{all-losses} L_j} \quad (1)$$

where δ_{obs} is the global average isotopic composition of CH_3Cl , δ_i and S_i are the isotopic composition and strength of source i , and ϵ_j and L_j are the isotopic fractionation effect and magnitude of sink j . Reported uncertainties have been calculated using Gaussian error propagation from the errors associated with the source compositions and sink fractionations mentioned above.

An important parameter currently unknown is the KIE for the reaction of CH_3Cl with OH-radicals (ϵ_{OH}). This value can be determined in laboratory studies and such measurements can be made with high accuracy (Cantrell et al. 1990; Rudolph et al. 2000). Thus we express the average isotopic composition as a function of the KIE for the loss due to OH-radical reaction by rearranging (1):

$$\frac{\sum_{all-sources} \delta_i * S_i}{\sum_{all-sources} S_i} = \delta_{obs} - \frac{\epsilon_{OH} L_{OH} + \sum_{NonOH-losses} \epsilon_j * L_j}{\sum_{all-losses} L_j} \quad (2)$$

The resulting average source composition (of all sources) can be described by -39.5‰ ($\pm 1.4\text{‰}$) $- 0.86 * \epsilon_{OH}$. Similarly, the average composition of sources with an unknown stable isotope ratio (including oceanic, coal combustion, incineration and the deficit with a total of 2.67 Tg $\text{CH}_3\text{Cl}/\text{yr}$) is -35.3‰ ($\pm 4.5\text{‰}$) $- 1.26 * \epsilon_{OH}$.

The current estimate for an oceanic source of CH_3Cl is 0.65 Tg $\text{CH}_3\text{Cl}/\text{yr}$ and we denote isotopic composition of these emissions as δ_{ocean} . This enables us to describe the stable isotopic composition of the deficit (δ_{def}) of 2.02 Tg $\text{CH}_3\text{Cl}/\text{yr}$ in the atmospheric CH_3Cl budget by the following expression:

$$\delta_{def} = -46.7\text{‰} (\pm 6.0\text{‰}) - 1.67 * \epsilon_{OH} - 0.32 * \delta_{ocean} \quad (3)$$

The deficit estimation includes a small contribution from minor sources with unknown stable carbon isotope ratios (Table 4), however it is extremely unlikely that these small emissions will significantly change the required stable carbon isotope ratio of the shortfall in the CH_3Cl budget. For example an isotopic composition of these small sources in the range between -80‰ and -5‰ will not result in a change of δ_{def} that exceeds its error of 6.0‰.

As previously mentioned, ϵ_{OH} is unknown. However, the KIE for the reaction of methane with OH radicals is positive (Cantrell et al. 1990), similar to recent observations of Rudolph et al. (2000) who found that the carbon isotope fractionation associated with the reaction of several alkanes, alkenes, and benzene with OH radicals results in an enrichment of ^{13}C in the remaining reservoir. Therefore it seems very likely that atmospheric removal of CH_3Cl by OH radicals will result in an enrichment of ^{13}C , although the magnitude of this effect is unknown. For the purposes of the following calculations, we have estimated the KIE (usually presented relative to unity as $\epsilon_{OH} = (KIE-1)*1000$ with units of ‰) to be $5 \pm 5\%$ (5‰ being the ϵ_{OH} for CH_4 and OH) although the true ϵ_{OH} could well be outside this range. Considering only loss via OH destruction, the required average CH_3Cl source composition is given by $\delta^{13}C_{observed} - \epsilon_{OH}$. Thus, using the measured global average CH_3Cl composition, the required average CH_3Cl source composition (over all sources) is $-41.2 \pm 5.0\%$. Including all known losses, the required average CH_3Cl source composition is $-43.8 \pm 4.5\%$.

The quantity with the lowest uncertainty in the budget of CH_3Cl is the atmospheric loss rate. We can separate the emissions required to balance the loss rate into three categories: those for which there are some indicators as to the isotopic composition (biomass burning, fungi, industry, and salt marsh emissions), those for which the composition is not known (oceanic, coal combustion, and incineration), and the deficit in the budget. This deficit may be due to errors in the strength of known sources or the existence of presently unidentified emissions. The weighted average of emissions from biomass burning, fungi, industry and salt marshes is $-48.7 \pm 8.8\%$ (1.23 Tg CH_3Cl/yr). This means, continuing the assumption that $\epsilon_{OH} = 5 \pm 5\%$, CH_3Cl emitted from all other sources must have an average isotopic composition of $-41.6 \pm 7.8\%$ (2.67 Tg CH_3Cl/yr).

Current understanding is that the large deficit in the atmospheric CH_3Cl budget is not due to underestimation of the magnitude of either the biomass burning source, or oceanic emissions. Lobert et al. 1999 constrained the upper limit of global biomass burning emissions to be 1.11 Tg CH_3Cl/yr . The ocean is insufficiently supersaturated in CH_3Cl to account for more than a small percentage of the total CH_3Cl budget (Moore et al. 1996). Clues as to unidentified sources include the results of Yokouchi et al. 2000, who found that levels of CH_3Cl in marine air samples were correlated with α -pinene, suggesting coastal terrestrial sources. Keppler et al. 2000 showed that alkylhalides could be produced abiotically from halide ions in organic matter (e.g. soil) via redox reactions, however the importance of this process is questionable and is reliant on the availability of the right source materials at sufficient concentration.

Since ϵ_{OH} will mostly likely be positive (see above), this budget indicates the unidentified sources of atmospheric CH_3Cl are relatively low in ^{13}C , that is significantly lower than carbon in fossil fuel and plant material. The results show that CH_3Cl emitted from the oceans would have to be very strongly depleted in ^{13}C (composition of around -90%) to allow the missing sources to have stable carbon isotope ratios similar to fossil fuel or terrestrial plant material. The required lightness, ie depletion in ^{13}C , of the missing source/s is compatible with what has been observed for emissions from biological sources and also processes such as biomass burning.

CFC113

The average tropospheric CFC113 isotopic composition was $-23.3 \pm 1.6\%$, $N = 38$ with no difference between the hemispheres. Since CFC113 has a lifetime of 85 years, and present day emissions are marginal due to the Montreal Protocol and its amendments, the global troposphere is well mixed with respect to CFC113. This means that the isotopic composition of tropospheric CFC113 will be homogenous and so the variability in the ambient observations is indicative only of the experimental uncertainty.

With only one recently produced commercial sample analysed, it is not justified to attach any significance to the difference in stable carbon isotope ratio between commercial and tropospheric CFC113. A substantial fractionation between emissions and ambient CFC113 is difficult to explain as CFC113 is unreactive. Also, the only significant tropospheric loss process is transport to the stratosphere. The fractionation due to stratospheric photolysis (the major loss mechanism in the stratosphere) of CFC113 is not known, however because of its long lifetime fractionation would need to be extremely large to cause any discernable change in the isotopic composition of tropospheric CFC113. There is nothing published on the stable carbon isotopic composition of CFC113 to date, however McCauley et al. (1999) found isotopic variability of up to 15% in commercial samples of bromomethane, and as mentioned above industrial CH_3Cl has been measured to be -60% and -41.9% .

Conclusions

The global average stable carbon isotopic composition of tropospheric CH_3Cl has a well defined average of $-36.2 \pm 0.3\%$. However further study is warranted. In particular, current Southern Hemispheric data are inadequate. The magnitude of the seasonal cycles (over a longer period and more

geographic coverage), and the North-South Hemispheric gradient need to be better defined.

The measurement of atmospheric CH_3Cl composition allows definition of the average isotopic composition of the unknown CH_3Cl source/s, as a function of the KIE for reaction with OH-radical. This average unknown source composition is depleted in ^{13}C relative to known potential parent materials (plants, fossil fuel) implying substantial fractionation of emission processes.

Aside from the obvious identification of the unknown sources, the key uncertainties in the isotopic budget of CH_3Cl are the KIE for loss via OH-radical and the isotopic composition of oceanic emissions. Also important is reducing the uncertainty in the isotopic composition of biomass burning emissions.

Acknowledgements

The authors would like to thank Gordon Brailsford, Janet Petersen, and Gavin Fisher from the National Institute of Water and Atmospheric Research of New Zealand (New Zealand samples), Jan Bottenheim, Doug Worthy (Fraserdale and Alert samples), and Darrell Ernst (technical assistance) from the Meteorological Service of Canada, Christine Rigby from Guelph University (Vancouver samples) and Larry Kleinman from Brookhaven National Laboratory (Houston samples). Thanks are also due to Markus Bill and Michael Gautois for the use of their unpublished data and helpful discussion. This work was funded by the National Science and Engineering Research Council of Canada. Alexandra Thompson thanks the American Association of University Women for an International Fellowship during this work.

References

- Andreae MO (1991) Biomass burning: its history, use and distribution and its impact on environmental quality and global climate. In: Levine JS (Ed) *Global Biomass Burning*. MIT Press, Cambridge, MA
- Bill M, Rhew RC, Weiss R & Goldstein A (2002) Carbon isotope ratios of methyl bromide and methyl chloride emitted from a coastal salt marsh. *Geophys. Res. Lett.* 29(4): 4-1-4-4, DOI 10.1029/2001GL012946
- Canirell CA, Shetter RE, McDaniel AH, Calvert JG, Davidson JA, Lowe DC, Tyler SC, Cicerone RJ & Greenberg JP (1990) Carbon kinetic isotope effect in the oxidation of methane by the hydroxyl radical. *J. Geophys. Res.* 95: 22455-22462
- von Czapiewski K, Czuba E, Huang L, Ernst D, Norman AL, Koppman R & Rudolph J (2002) Isotopic composition of non methane hydrocarbons in emissions from biomass burning. *Atmos. Chem.* (in press)

- Gautrois M, Brauers TH, Koppmann R, Rohrer F, Stein O & Rudolph J (Manuscript in preparation) Seasonal variation and trends of volatile organic compounds in the lower polar troposphere (pers. comm.)
- Harper DB, Kalin RM, Hamilton JTG & Lamb C (2001) Carbon isotope ratios for chloromethane of biological origin: potential tool in determining biological emissions. *Environ. Sci. Tech.* 35: 3616–3619
- Intergovernmental Panel on Climate Change (IPCC). *Climate Change 1995: Radiative Forcing of Climate Change and an Evaluation of the IPCC IS92 Emission Scenarios*. Cambridge University Press, Cambridge, U.K.
- Kaplan IR (1975) Stable isotopes as a guide to biogeochemical processes. *Proc. Roy. Soc. London, Series B* 189: 183–211
- Keene WC, Khalil AK, Erickson III DJ, McCulloch A, Graedel TE, Lobert JM, Aucott ML, Gong SL, Harper DB, Kleiman G, Midgley P, Moore RM, Seuzaret C, Sturges WT, Benkovitz CM, Koropalov V, Barrie LA & Li YF (1999) Composite global emissions of reactive chlorine from anthropogenic and natural sources: Reactive Chlorine Emissions Inventory. *J. Geophys. Res.* 104: 8429–8440
- Kepler F, Eiden R, Niedan V, Pracht J & Schoeler HF (2000) Halocarbons produced by natural oxidation processes during degradation of organic matter. *Nature* 403: 298–301
- Khalil MAK, Moore RM, Harper DB, Lobert JM, Erickson DJ, Koropalov V, Sturges WT & Keene WC (1999) Natural emissions of chlorine-containing gases: Reactive Emissions Inventory. *J. Geophys. Res.* 104: 8333–8346
- Khalil MAK & Rasmussen RA (1999) Atmospheric methyl chloride. *Atmos. Environ.* 33: 1305–1321
- Lobert JM, Keene WC, Logan JA & Yevich R (1999) Global chlorine emissions from biomass burning: Reactive Chlorine Emissions Inventory. *J. Geophys. Res.* 104: 8373–8389
- Lowe DC, Brenninkmeijer CAM, Brailsford GW, Lassey KR & Gomez AJ (1994) Concentration and ¹³C records of atmospheric methane in New Zealand Antarctica: Evidence for changes in methane sources. *J. Geophys. Res.* 99: 16913–16925
- Matthews DE & Hayes JM (1978) Isotope ratio monitoring gas chromatography-mass spectrometry. *Anal. Chem.* 50: 1465–1473
- McCauley SE, Goldstein AH & DePaolo DJ (1999) An isotopic approach for understanding the CH₃Br budget of the atmosphere. *Proc. Natl. Acad. Sci. U.S.A.* 96: 10006–10009
- Miller LG, Kalin RM, McCauley SE, Hamilton JTG, Harper DB, Millet DB, Oremland RS & Goldstein AH (2001) Large carbon isotope fractionation associated with oxidation of methyl halides by methylotrophic bacteria. *Proc. Nat. Acad. Sci. U.S.A.* 98: 5833–5837
- Mook WG, Koopmans M, Carter AF & Keeling CD (1983) Seasonal latitudinal and secular variations in the abundance and isotope ratios of atmospheric carbon dioxide. *J. Geophys. Res.* 88: 10915–10933
- Moore RM, Groszko W & Niven S (1996) Ocean – atmosphere exchange of methyl chloride: results from NW Atlantic and Pacific Ocean studies. *J. Geophys. Res.* 101: 28529–28539
- Rhew RC, Miller B & Weiss R (2000) Natural methyl bromide and methyl chloride emissions from coastal salt marshes. *Nature* 403: 292–295
- Rudolph J, Lowe D, Martin RJ & Clarkson TS (1997) A novel method for compound specific determination of $\delta^{13}\text{C}$ in volatile organic compounds at ppt levels in ambient air. *Geophys. Res. Lett.* 24: 659–662
- Rudolph J, Czuba E & Huang L (2000) The stable carbon isotope fractionation for reactions of selected hydrocarbons with OH-radicals and its relevance for atmospheric chemistry. *J. Geophys. Res.* 105: 29329–29346

Tsunogai U, Yoshida N & Gamo T (1999) Carbon isotopic compositions of C2–C5 hydrocarbons and methyl chloride in urban, coastal, and maritime atmospheres over the western North Pacific. *J. Geophys. Res.* 104: 16033–16039

Yokouchi Y, Toom-Saunry D & Barrie LA (1998) Measurements of methyl halides (CH₃Cl, CH₃Br, CH₃I) at Alert. Canadian Baseline Program: Summary of Progress to 1998

Yokouchi Y, Nojiri Y, Barrie L, Toom-Saunry D, Machida T, Inuzuka Y, Akimoto H, Li H-J, Fujinuma Y & Aoki S (2000) A strong source of methyl chloride to the atmosphere from tropical coastal land. *Nature* 403: 295–298

