

## Centre – The Middle of a Distribution

- ▶ An average is the single value that best represents the centre of a series of values.
- ▶ These values could be the shooting percentage of your favourite basketball team or the average rate at which convicted criminals re-offend.
- ▶ An average is like the fulcrum of a seesaw – the point at which a seesaw is perfectly level.



“Seesaw” by dianaholga: CC BY-NC-ND 2.0

## Centre – The Middle of a Distribution

- ▶ Averages are known as **measures of central tendency** and have three variations, the
  1. Mean
  2. Median
  3. Mode




"Seesaw" by dianaholga: CC BY-NC-ND 2.0

## The Mean

$$\bar{X} = \frac{\sum X_i}{n}$$

- ▶  $\bar{X}$  is the mean.
- ▶  $\sum$ , the Greek letter sigma, instructs us to sum or add-up.
- ▶  $X_i$  is the individual value.
- ▶  $n$  is the number of values in a series, often called the **sample** or **population size**.

Parking tickets issued by the City of Guelph, Ontario



## The Mean – Maple Leafs Player Salaries 2021-22



### Stem-and-Leaf Plot

0	77789999
1	0225566
2	05
3	58
4	
5	006
6	
7	0
8	
9	
10	9
11	06

Mean = \$3.23 million

## The Mean

$$\bar{X} = \frac{\sum X_i}{n}$$

- ▶ The Mean is the Average or most central score.
- ▶ Sometimes a capital  $N$  represents the number of values. Other times, it is represented by a small  $n$ .

- ▶ The mean is like a fulcrum on a seesaw – the weight of the values below the mean equals the weight of the values above of the mean.



“Seesaw” by dianaholga: CC BY-NC-ND 2.0

## The Mean – Maple Leafs Player Salaries 2021-22



### Stem-and-Leaf Plot

0	77789999
1	0225566
2	05
3	58
4	
5	006
6	
7	0
8	
9	
10	9
11	06


Mean = \$3.23 million

## The Mean

$$\bar{X} = \frac{\sum X_i}{n}$$

- ▶ A strength and weakness of the mean is its sensitive to extreme values (also known as **outliers**).

### The Toronto Raptors Salary: 2021-22




Mean = \$8.5 million

## The Mean

$$\bar{X} = \frac{\sum X_i}{n}$$

- ▶ A strength and weakness of the mean is its sensitive to extreme values (also known as **outliers**).

## The Toronto Raptors Salary: 2021-22



Mean without Pascal Siakam = \$5.67 million



## The Mean

Five Largest Toronto Raptors Salaries  
2021-22 (rounded) – Stem and Leaf  
Diagram

1		669
2		0
3		3

*in tens of millions of US dollars*

- ▶ Find the mean salary for the five NBA players.
- ▶ If we added a sixth NBA player, what salary would he need to have for the mean to become \$16 million?

## The Mean

$$\bar{X} = \frac{\sum X_i}{n}$$

- ▶ The **arithmetic mean**: the sum of the deviations from the mean is equal to zero.

Deviation from the Mean =  $X_i - \bar{X}$

### Salaries

Player 1	4,000,000
Player 2	5,000,000
Player 3	6,000,000

## The Weighted Mean

How to Compute a Weighted Mean

- ▶ Find the value by multiplying the frequency and the parking fine.

Parking Tickets – City of Guelph

Fine	Frequency	Value
\$30	2,209	
\$35	13,007	
\$40	5,410	
\$51	1,271	
\$60	1,488	
\$91	2,132	

Find the weight mean fine on a parking ticket.

## The Weighted Mean

How to Compute a Weighted Mean

- ▶ Find the value by multiplying the frequency and the parking fine.

Parking Tickets – City of Guelph

Fine	Frequency	Value
\$30	2,209	66,270
\$35	13,007	455,245
\$40	5,410	216,400
\$51	1,271	64,821
\$60	1,488	89,280
\$91	2,132	194,012

Find the weight mean fine on a parking ticket.

## The Weighted Mean

How to Compute a Weighted Mean

- ▶ Find the value by multiplying the frequency and the parking fine.
- ▶ Sum all the values and frequencies.

Parking Tickets – City of Guelph

Fine	Frequency	Value
\$30	2,209	66,270
\$35	13,007	455,245
\$40	5,410	216,400
\$51	1,271	64,821
\$60	1,488	89,280
\$91	2,132	194,012

Find the weight mean fine on a parking ticket.

## The Weighted Mean

How to Compute a Weighted Mean

- ▶ Find the value by multiplying the frequency and the parking fine.
- ▶ Sum all the values and frequencies.

Parking Tickets – City of Guelph

Fine	Frequency	Value
\$30	2,209	66,270
\$35	13,007	455,245
\$40	5,410	216,400
\$51	1,271	64,821
\$60	1,488	89,280
\$91	2,132	194,012
Totals	25517	1086028

Find the weight mean fine on a parking ticket.

## The Weighted Mean

### How to Compute a Weighted Mean

- ▶ Find the value by multiplying the frequency and the parking fine.
- ▶ Sum all the values and frequencies.
- ▶ Divide the values by the total frequency to find the weighted average.

Parking Tickets – City of Guelph

Fine	Frequency	Value
\$30	2,209	66,270
\$35	13,007	455,245
\$40	5,410	216,400
\$51	1,271	64,821
\$60	1,488	89,280
\$91	2,132	194,012
	25,517	1,086,028

## The Weighted Mean

### How to Compute a Weighted Mean

- ▶ Find the value by multiplying the frequency and the parking fine.
- ▶ Sum all the values and frequencies.
- ▶ Divide the values by the total frequency to find the weighted average.

Parking Tickets – City of Guelph

Fine	Frequency	Value
\$30	2,209	66,270
\$35	13,007	455,245
\$40	5,410	216,400
\$51	1,271	64,821
\$60	1,488	89,280
\$91	2,132	194,012
	25,517	1,086,028

Find the weight mean fine on a parking ticket:  
 $1,086,028 / 25,517 = \$42.6$



## The Weighted Mean: Weight on Each Fine Dollar Amount

Parking Tickets – City of Guelph

Fine	Frequency	Value	Weight
\$30	2,209	66,270	
\$35	13,007	455,245	
\$40	5,410	216,400	
\$51	1,271	64,821	
\$60	1,488	89,280	
\$91	2,132	194,012	
	25,517	1,086,028	

- ▶ Find the weighting on each ticket fine by divide the value of each fine by the total value.

## The Weighted Mean

- Find the weighting on each ticket fine by divide the value of each fine by the total value.

Parking Tickets – City of Guelph

Fine	Frequency	Value	Weight
\$30	2,209	66,270	6.1%
\$35	13,007	455,245	41.9%
\$40	5,410	216,400	19.9%
\$51	1,271	64,821	6.0%
\$60	1,488	89,280	8.2%
\$91	2,132	194,012	17.9%
	25,517	1,086,028	100%

*The weight represents the relative contribution of each fine to total revenue collected.*

## The Median

The median is

- ▶ a different kind of **average**.
- ▶ the **midpoint** of a series where half the values are below the median and half the values are above the median.

To find the median:

- ▶ List the values in order.
- ▶ Find the middle score.
- ▶ If there are an even number of values, the two middle values are the median.

Maple Leafs Salaries 2021-22 –  
Stem-and-Leaf Plot

0	77789999
1	0225566
2	05
3	58
4	
5	006
6	
7	0
8	
9	
10	9
11	06

## The Median

The median is

- ▶ a different kind of average.
- ▶ the **midpoint** of a series where half the values are below the median and half the values are above the median.

To find the median:

- ▶ List the values in order.
- ▶ Find the middle score. ]item If there are an even number of values, the two middle values are the median.

Five Largest Toronto Raptors Salaries  
2021-22 (rounded) – Stem and Leaf

1		669
2		0
3		3

## The Median v Mean


$$X = \{2, 7, 9\}$$

- ▶ The mean is the middle point where
  - ▶ the value below and above the mean are equally weighted, and
  - ▶ the sum of the deviations from the mean equals zero.
- ▶ The median is the middle point of a set of cases.
- ▶ The median is insensitive to extreme values, while the mean is sensitive to extreme values.

## The Median v Mean


- ▶ The mean is the middle point where
  - ▶ the value below and above the mean are equally weighted, and
  - ▶ the sum of the deviations from the mean equals zero.
- ▶ The median is the middle point of a set of cases.
- ▶ The median is insensitive to extreme values, while the mean is sensitive to extreme values.

The Toronto Raptors Salary: 2021-22



Mean = \$8.5 million  
Median = \$2.0 million

## The Median v Mean



## Stem-and-Leaf Plot for Maple Leafs


Salaries 2021-22

```
0 | 77789999
1 | 0225566
2 | 05
3 | 58
4 |
5 | 006
6 |
7 | 0
8 |
9 |
10 | 9
11 | 06
```

Mean = \$3.23 million  
Median = \$1.55 million


## The Mode

- ▶ The mode is the value that occurs most often.
- ▶ Of the four crimes, which occurs most often?





## What is the Mode?



### Toronto Maple Leafs Salaries 2021-22

0	77789999
1	0225566
2	05
3	58
4	
5	006
6	
7	0
8	
9	
10	9
11	06

Mean: \$3.23 million

Median: \$1.55 million

Mode:

Find the Mode Response

Student Responses to a Test Question

	A	B	C	D	E
Frequency	22	31	44	45	9

## What is Variability?

- ▶ **Variability** reflects the difference in values.
- ▶ Variability is also known as **dispersion** or **spread**.  $X = \{3, 3, 4, 4\}$  and  $Y = \{0, 1, 6, 7\}$
- ▶ Compute the Mean of  $X$  and  $Y$ :
- ▶ Compute the Deviation from the Mean for  $X$  and  $Y$ .

## What is Variability?

$X = \{3, 3, 4, 4\}$  and  $Y = \{0, 1, 6, 7\}$   
and  $Z = \{3.5, 3.5, 3.5, 3.5\}$

- ▶ Do any values in  $Z$  deviate from the mean?
- ▶  $Z$  has no variability at all!

## Three measures of variability

- ▶ The Range
- ▶ The Variance
- ▶ The Standard Deviation



"Ruby dice" by TomNatt is licensed under CC BY-NC 2.0

## The Range

- ▶ The range measures how far apart the values are.
- ▶ It is measured as the highest value in a series less the lowest value in a series.

Find the range for  $X$ ,  $Y$ ,  $Z$ :

$$X = \{3, 3, 4, 4\}$$

$$Y = \{0, 1, 7, 8\}$$

$$Z = \{3.5, 3.5, 3.5, 3.5\}$$

## Find the Range


### Toronto Maple Leafs Salaries 2021-22

0	77789999
1	0225566
2	05
3	58
4	
5	006
6	
7	0
8	
9	
10	9
11	06

### Selected Toronto Raptors Salaries 2021-22

0.	06,07,07
1.	669
2.	0
3*	3


## Find the range





## The Range


- ▶ The range is simple and only provides information about the end points which might be extremely large or small.
- ▶ The range provides no information on the interior values.
- ▶ Standard Deviation and Variance are more common measures of the spread (dispersion).



## The Interquartile Range (IQR)

- ▶ The interquartile range ignores the most extreme values and focuses on the middle of the data.
- ▶ Remember, the median divided the set of values into two halves: one above and one below the median.
- ▶ The interquartile range divides the set of values into four quarters.
- ▶ One quarter of the values in the set are below the lower quartile and one quarter of the values in the set are above the upper quartile.


# Find the Interquartile Range (IQR)



## How to Find the Interquartile Range (IQR)


### How to Find the Interquartile Range (IQR)

- ▶ Find the median of all values in the set.
- ▶ Find the median of the lower half of values (Q1).
- ▶ Find the median of the upper half of values (Q3).



## Why the Quartile Range?

- ▶ Finding the IQR and the range give more information about the distribution of values than just the range.
- ▶ Sometimes, researchers study quintiles (5 equal parts) or deciles (10 equal parts).
- ▶ Software programs can easily find these for you.



## The Standard Deviation is:

- ▶ A measure of *average variability* in a set of values.
- ▶ The average distance from the mean.

If the standard deviation is larger, the set of values have more variability.

$$\sigma = \sqrt{\frac{\sum (X_i - \bar{X})^2}{n}}$$

- ▶  $\sigma$  is the standard deviation.
- ▶  $\Sigma$  is sigma, the command to sum what follows.
- ▶  $X_i$  represents each individual value.
- ▶  $X_i - \bar{X}$  is the difference between an individual value and the mean.

Find the standard deviations:

$$X = \{3, 3, 5, 5\}$$

$$Y = \{1, 1, 7, 7\}$$


$$Z = \{4, 4, 4, 4\}$$

$$\sigma = \sqrt{\frac{\Sigma (X_i - \bar{X})^2}{n}}$$

- ▶  $\sigma$  is the standard deviation.
- ▶  $\Sigma$  is sigma, the command to sum what follows.
- ▶  $X_i$  represents each individual value.
- ▶  $X_i - \bar{X}$  is the difference between an individual value and the mean.

## How to Find the Standard Deviation

- ▶ Compute the mean
- ▶ Subtract the mean from each score
- ▶ Square each individual difference.
- ▶ Sum all the squared deviations.
- ▶ Divide the sum by  $n$ .
- ▶ Take the square root of the sum divided by  $n$ .





## The Standard Deviation

Calgary to:	Distance	Deviation	Squared
Winnipeg	1327 km		
Bismarck	1286 km		
Vancouver	1057 km		

## The Standard Deviation

Calgary to:	Kilometers	Deviation	Squared
Winnipeg	1327	103.7	10746.8
Bismarck	1286	62.7	3927.1
Vancouver	1057	-166.3	27666.8
	Mean	1223.3	
	Sum		42340.7
	Sum $\div$ 3		14113.6
	Squared-Root		118.8

## The Standard Deviation

Calgary to:	Kilometers	Deviation	Squared
Winnipeg	1327	103.7	10746.8
Bismarck	1286	62.7	3927.1
Vancouver	1057	-166.3	27666.8
	Mean	1223.3	
	Sum		42340.7
	Sum $\div$ 3		14113.6
	Squared-Root		118.8

Why square each value?

- ▶ Squaring each value removes negatives.
- ▶ If we simply added up the deviations from the mean, the sum would equal zero.

Why take the square root of the sum of squared-values?

- ▶ The square root returns the values into the same units from which they started.

## Statistical Bias

$$\sigma = \sqrt{\frac{\sum (X_i - \bar{X})^2}{n}}$$

- ▶  $\sigma$  is useful for summarizing information about a population.
- ▶ However if you want to make a statistical inference about a population using a sample,  $\sigma$  is a **statistically biased** estimate of the variability in a population.
- ▶  $\sigma$  *underestimates* the true variability in a population.

- ▶ For making statistical inferences, the **sample standard deviation**,  $s$ , is unbiased.

$$s = \sqrt{\frac{\sum (X_i - \bar{X})^2}{n - 1}}$$

- ▶ where the denominator is now  $n - 1$ .
- ▶ Dividing by a smaller value,  $n - 1$ , makes the standard deviation larger.
- ▶ Allowing for a larger standard deviation is a *conservative* estimate.

## Statistical Bias

$$s = \sqrt{\frac{\sum (X_i - \bar{X})^2}{n-1}} \quad \sigma = \sqrt{\frac{\sum (X_i - \bar{X})^2}{n}}$$

Sample Size	$\sum (X_i - \bar{X})^2$	$\sigma$	$s$	$s - \sigma$
10	600	7.75	8.16	0.4190
100	600	2.45	2.46	0.0123
1000	600	0.77	0.77	0.0004

- ▶ All else the same, as the sample size increases the difference between the standard deviation and the sample standard deviation becomes smaller.

## The Standard Deviation

- ▶ If the standard deviation is larger, the values are more spread out.
- ▶ If the standard deviation is larger, the values are more different from each other.
- ▶ The standard deviation is sensitive to extremely large or small values.
- ▶ An extremely large or small value is one that is far away from the mean!

## The Variance

- ▶ The variance is closely related to the standard deviation.
- ▶ The variance is equal to the standard deviation squared.

The Variance is  $\sigma^2 = \frac{\sum (X_i - \bar{X})^2}{n}$

A Previous Example:

Calgary to:	Kilometers	Deviation	Squared
Winnipeg	1327	103.7	10746.8
Bismarck	1286	62.7	3927.1
Vancouver	1057	-166.3	27666.8
	Mean	1223.3	
	Sum		42340.7
	Sum ÷ 3		14113.6
	Squared-Root		118.8


## The Variance

- ▶ Interpreting the variance can be more difficult than interpreting the standard deviation which is stated in the same units as the series.
- ▶ So,
  - ▶ If the values are stated in kilometers, the standard deviation is also stated in kilometers.
  - ▶ If the values are stated in dollars, the standard deviation is also stated in dollars.
- ▶ However, variance does have useful applications in finance, economics, and statistics for the social sciences.





## Data as a Distribution: Probability Density Function (PDF)

- ▶ Data can be represented according to frequency, a **frequency distribution**.
- ▶ If the data is continuous, values can be grouped into **class intervals**.
- ▶ A **class interval** is a range of values (numbers).
- ▶ Here, the class interval for soccer player heights is one-inch, for a total of 14 class intervals: 60-61, 61-62, ..., 73-74.




## Data as a Distribution: Probability Density Function (PDF)




# Data as a Distribution: Cumulative Density Function (CDF)

## PDF




## CDF



# Cumulative Frequency & Cumulative Probability

Range	Frequency	Cumulative Frequency	Probability	Cumulative Probability





## CDF of a Soccer Player Heights With 0.5 Inch Intervals

When working with continuous numeric data,

1. Sort the data from lowest to highest.
2. Divide the Data (values) into bins.
3. Graph the result.

