

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

Facial expressions during infancy are important to examine as infants do not have the language skills to describe their experiences. This is particularly vital in the context of pain where infants depend solely on their caregivers for relief. The objective of the current study was to investigate the development of negative infant facial expressions in response to immunization pain over the first year of life. Infant facial expressions were examined longitudinally using a subsample of 100 infants that were each videotaped during their 2-, 4-, 6-, and 12-month routine immunization appointments. Infant facial expressions were coded using BabyFACS for the first minute after a painful needle prick. Facial expressions were examined with a catalogue of the most commonly occurring facial expressions. Results demonstrated that clear differences were seen over ages. Infants display a variety of facial expressions with some of the components of adult pain expressions immediately after the needle and abate shortly after. However, infants did not display adult expressions of discrete negative emotions. Instead, infants display a variety of generalized pain and distress faces aimed at gaining caregiver aid. The development of non-verbal communication in infants, particular facial expressions, remains an important area of inquiry. Further study into accurately measuring infant negative emotions, pain, and distress is warranted.

Keywords: immunization, infant, facial expression, emotion, acute pain

**A Longitudinal Analysis of the Development of Infant Facial Expressions in
Response to Acute Pain: Immediate and Regulatory Expressions**

Sara Ahola Kohut¹

Rebecca Pillai Riddell^{1,2,3}

David B. Flora¹

Harriet Oster⁴

¹York University, Toronto, Canada

²Hospital for Sick Children, Toronto, Canada

³University of Toronto, Toronto, Canada

⁴New York University, New York, U.S.A.

Corresponding Author:

Dr. Rebecca Pillai Riddell, Ph.D., C.Psych.

The Opportunities to Understand Childhood Hurt (OUCH) Laboratory

York University, Department of Psychology, Faculty of Health

119 Behavioural Sciences Building

4700 Keele Street, Toronto, Ontario, Canada, M3J 1P3

Telephone: 416-736-2011 ext 20177, Fax: 416-736-5814

E-mail: rpr@yorku.ca, Website: www.yorku.ca/ouchlab/

31 pages in total (3 tables and 6 figures)

Introduction

The development of negative emotions in a pain context is an important area of study because the underpinnings of regulation from distress (pain-related or otherwise) occur in the first year of life [4]. Researchers have been divided between two major perspectives. According to Differential Emotions Theory, infants show clearly differentiated facial expressions of discrete negative emotions such as anger, fear, and pain from early infancy [16,18]. On the other hand, according to classic differentiation theories, facial expressions of specific, discrete negative emotions gradually differentiate from more generalized, global distress expressions [6,8,37]. In this vein, Oster's more recent Ontogenetic perspective holds that infant facial and vocal expressions are neither precocious, immature adult expressions nor global and diffuse signs of arousal. Instead, they are distinctive behavioral adaptations that evolved because they serve crucial communicative functions in infancy [24].

The current study contributes to our knowledge of the development of negatively-valenced facial expressions and emotions related to pain-related distress by following a group of infants undergoing routine immunizations. Immunizations allow us to examine infant facial expressions in response to a distressing stimulus in an ecologically valid setting. However, previous work studying presumed discrete emotions in the immunization context is limited by the use of coding systems that assume adult emotional prototypes and small sample sizes [19,20].

While we expected that we would observe a distinct pain face (validly and reliably measured by the well-established Neonatal Facial Coding System [14,15]), we

1
2
3
4
5
6 did not limit our scope to only a pain face. Thus, if there were non-pain facial expressions
7
8 present (e.g., anger, fear) or variations of pain faces present, we wanted to be able to
9
10 capture these constellations and their development over the first year of life. Only the
11
12 Facial Action Coding System for Infants and Young Children (BabyFACS [23]) codes
13
14 the full range of anatomical facial actions possible. Utilizing a comprehensive coding
15
16 system, one that does not only code facial actions present in pain expressions but rather
17
18 the whole range of emotional facial displays, allows the opportunity to better understand
19
20 the infant's emotional experience within the acute pain context. To our knowledge, this is
21
22 the only extant study that uses this exhaustive system to track the development of facial
23
24 expressions over the immunization appointment (i.e. for one minute post needle) and over
25
26 age (i.e. at 2, 4, 6 and 12 months of age).
27
28
29
30
31

32
33 We aim to address three questions: 1) What are the most commonly occurring
34
35 facial expressions after an acutely painful stimulus during the first year of life? 2) Does
36
37 the average time spent in these different expressions change over infant age? 3) How
38
39 does the occurrence of the commonly occurring facial expression change as infants
40
41 regulate distress over the first minute after painful stimulus? We hypothesized that, in
42
43 line with a differentiation model of emotions, pain-elicited facial expressions become
44
45 more differentiated and complex with age. We also hypothesized, specifically in line
46
47 with the Ontogenetic perspective, that infant-specific facial expressions, distinct from
48
49 prototypical adult expressions, would be found.
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Methods

Participants

One hundred infants were randomly selected from a large cohort of infants being observed during four separate routine immunization appointments over the first year of life [27]. At the time of data analysis, 598 infants had been recruited into the cohort with 131 infants having some video footage at all four appointments. Infants weighed on average 7.43 lbs ($SD = 0.92$) at birth, and were most often the first or second born child in the family (52% 1st born and 42% 2nd born). Half of the sample was male (50%). Mothers were most often the primary caregiver and had an average age of 34.76 years ($SD = 4.82$). Mothers were most often married (87%) and educated (74% minimum university educated). Infants were also culturally diverse with 83% of mothers reporting a heritage culture other than North American. Infants at the 2-, 4-, and 6-month immunization appointments most often received a Pediacel vaccine first, followed by a Pentacel vaccine. At the 12-month appointment, infants received a Pentacel vaccine and either a MMR vaccine or a Priorix vaccine. Refer to Table 1 for a detailed summary of demographic information.

Procedure

Ethical approval was obtained through research ethics review boards at both the participating university and the associated pediatric hospital. Briefly, healthy infants were observed during routine immunization appointments at 2, 4, 6, and 12 months of age. During these appointments, infants' facial expressions were video recorded before and after the immunization. Caregivers completed demographic and health questionnaires at

each visit. A detailed description of the procedures for the larger study is previously described and available for review [27].

Facial Coding Measure

Infant facial expressions were coded for one minute after the last immunization needle using BabyFACS [23]. BabyFACS is an adaptation for infants and young children of the adult Facial Action Coding System (FACS) [10,11] that accounts for the differences in infant facial morphology. It is a comprehensive anatomically based measure of all visually discernable facial movements in infants. BabyFACS does not provide formulas for facial expressions of emotions but rather provides a method to describe constellations of facial actions displayed without *a priori* assumptions of what facial actions may be displayed based on adult facial expressions. BabyFACS includes 73 facial muscle actions (each identified by a number) coded for intensity from A (trace) to E (maximum). This coding creates full-face scores that describe precisely which facial muscles are activated and how strongly they are activated. BabyFACS also includes 3-by-3 matrices for coding intensities and variants of smiles and cry faces. BabyFACS coding allows for descriptions of infant facial expressions without prior assumptions of their meaning and therefore provides an objective measure of infant facial expressions.

For the current analysis, coding was done in two stages. First, 100% of the footage was double-coded by certified BabyFACS coders to ensure an agreed upon time for the onset, apex and offset of each facial expression. Second, using these agreed onset, apex and offset times, footage was independently coded by the two coders on BabyFACS. Twenty-five percent of the data was double-coded for inter-rater reliability

and, based on the FACS and BabyFACS manuals method, the reliability coefficient was deemed acceptable at 0.84.

Coding and Analysis Procedure

Infant facial expressions were coded for one minute after the final immunization needle of the appointment. Categories of facial expressions were determined in a stepwise manner led by the lead author in collaboration with a coauthor and then independently confirmed with the author of BabyFACS. Categories were explored via frequency of facial expressions across the sample of infants as well as via duration of the facial expressions. Initial steps in creating categories included an examination of facial actions that might be able to cluster together due to anatomic similarity. In the eyes, AU6 results in the raising of the cheeks and skin below the eyes, and AU7 results in squinting and tightening the lower eyelids; both narrow the eye opening. These two facial actions (AU6 and AU7) are difficult to distinguish in infants within the pain context and were clustered together. In the midface, AU9, AU10, and AU11 all deepen the nasolabial furrow and raise the upper lip; these facial actions often co-occur and can be difficult to distinguish in distress expressions. These three facial actions (AU9, AU10, and AU11) were therefore clustered together. In the mouth, AU26 and AU27 are related to how widely a mouth is vertically opened. In the context of a cry face, these facial actions both reflect a dropping of the jaw and were clustered together. A subsequent review of the adult literature on facial expressions of pain confirmed that these clusters of facial actions were appropriate given our context [34]. Next, based on the potential conceptual importance of certain facial actions and sub-constellations (namely AU43, eyes closed;

1
2
3
4
5
6 AU6 or AU7, cheek raise or tightening of the eyelid; AU20, horizontal stretch; cry mouth
7
8 matrix [22,33]), the categories were further refined. Categories were collapsed and
9
10 adapted to create fewer groupings with consideration of particular facial actions and cry
11
12 mouth combinations. Seven categories of facial expression constellations of negative
13
14 affect were agreed upon based on frequency, inclusivity, mutual exclusiveness, and
15
16 observable differences of categories.
17
18

19
20 To avoid *a priori* assumptions regarding the interpretation of facial expressions
21
22 during our initial coding and synthesis, the most commonly occurring facial
23
24 configurations were randomly assigned a colour name. To answer our first question, we
25
26 first describe a catalogue of each of seven discerned facial expressions (Red, Orange,
27
28 Yellow, Green, Blue, Purple, and Black) along with the mean percentage of time over the
29
30 minute that infants spent displaying that facial expression. To answer our second
31
32 question, statistical analyses compared the average percent of the minute each facial
33
34 expressions was displayed across age (2-, 4-, 6-, and 12-months) using a related-samples
35
36 Friedman's two-way ANOVA by ranks for each facial expression. A Bonferonni
37
38 correction was applied such that $p < 0.001$ was required for significance. Significant
39
40 ANOVAs were followed by simple pairwise comparisons. Finally, to answer the third
41
42 question, the 1-minute post-needle epoch was broken up into 5-second increments and
43
44 the average percentage of time within each 5-second sub-epoch was graphed to describe
45
46 how the occurrence of facial displays changed over the first minute after the last needle.
47
48
49
50
51
52

53
54 For discussion and interpretation of our findings, each general category (facial
55
56 expression) was also coded based on the Neonatal Facial Coding System (NFCS [14,15]),
57
58
59
60
61
62
63
64
65

The Maximally Discriminative Facial Movement coding system (Max [17]), BabyFACS [23], and an adult pain metric [32,35] based on FACS Action Units [10,11]. NFCS pain scores were calculated by a certified NFCS coder based on presence or absence of seven facial actions (brow bulge, eye squeeze, nasolabial furrow, open lips, horizontal stretch mouth, vertical stretch mouth, taut tongue) with higher scores indicative of higher pain on a scale from 0 to 7. Max scoring was done by a Max-certified coder based on emotion labels defined in the manual. BabyFACS interpretations were determined based on the manual by the creator of the measure. FACS AUs were coded based on the manual by a certified FACS coder. The Adult pain metric [32, 35] was calculated based on the sum of the intensities of AU4 + AU6 or AU7 +AU9 or AU10 + AU43. All AUs are coded based on intensity (A to E which translates to a score 0 to 5) except AU43 which is coded present or absent (0 or 1) based on the maximum intensity, AU 43E (eyes closed). Higher scores are indicative of higher pain and on a scale of 0 to 16. However, since scoring this pain metric for this study was calculated for a general category of a facial expression, intensity scores of AUs were based on the most common intensity scores within the category (C to D which translates to a score of 3 or 4). Therefore, the possible range of scores was 0 to 13 [32,35].

Results

The seven facial configurations identified are described in Table 2 and Figure 1. Mouth descriptions (vertical and horizontal stretching) are based on the cry matrix described in the BabyFACS manual [23]. Analyses are described according to each facial expression (i.e. colour label), alongside a pictorial example.

1
2
3
4
5
6 *Red* (see Figure 2): Overall average percentage of Red facial expression
7
8 (collapsed over the minute post-needle) were significantly different across age ($\chi^2 (3) =$
9 34.58, $p < .001$; 2-month: 7.49%, 4-month: 5.12%, 6-month: 9.51%, and 12-month:
10 18.87%). Post hoc pairwise comparisons indicated a significant increase in Red facial
11 expressions in infants from 2 to 12 months and from 4 to 12 months. Next, the trajectory
12 of average percentage per 5-second increments over the minute for Red facial expressions
13 was graphed separately for each age. In the first 30 seconds immediately after the last
14 immunization needle, for each age there were increasing displays of Red facial
15 expressions. In the 30 to 59 seconds following the last immunization needle, for each age
16 there were decreasing displays of Red facial expressions over the minute. Despite their
17 opposing direction, the amount of change during the first 30 seconds appears similar to
18 that for the last 30 seconds. Finally, infants at 12 months displayed more Red facial
19 expressions than at 2, 4, and 6 months across the entire minute.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

37 *Orange*: The overall average percentage of Orange facial expressions occurred
38 less than 5% of the minute following the needle with no significant difference across age
39 ($\chi^2 (3) = 6.61$, $p = .09$; 2-month: 1.36%, 4-month: 3.07%, 6-month: 2.44% and 12-month:
40 2.37%). Overall, infants consistently displayed Orange facial expressions less than 5% of
41 the time during every 5-second increment of the minute.
42
43
44
45
46
47
48
49

50 *Yellow* (see Figure 3): The average percentage of Yellow facial expressions over
51 the minute remained generally stable across age ($\chi^2 (3) = 10.09$, $p = .02$; 2-month:
52 17.09%, 4-month: 10.36%, 6-month: 9.63% and 12-month: 13.46%). In the first 15
53 seconds, for each age there were increasing displays of Yellow facial expressions. Infants
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6 at 12 months displayed the most Yellow facial expressions during the first 10 seconds. In
7
8 the last 15 to 59 seconds, there was a decrease in the occurrence of Yellow facial
9
10 expressions. During this later period, 2-month-olds continued to show high levels of
11
12 Yellow expressions, while the levels dropped sharply at the 12-month appointment.
13
14

15 *Green* (see Figure 4): There was a significant difference by age in overall mean
16 occurrence over the minute ($\chi^2(3) = 16.96, p < .001$; 2-month: 2.49%, 4-month: 2.40%,
17
18 6-month: 2.51% and 12-month: 0.00%), but no pairwise comparisons were significant. At
19
20 12 months, infants did not display Green facial expressions in response to the last
21
22 immunization needle. For infants at 2, 4, and 6 months, Green facial expressions occur
23
24 within the first 10 seconds after immunization needle then diminish until the end of the
25
26 one minute epoch.
27
28
29
30
31

32 *Blue* (see Figure 5): Blue facial expressions significantly differed in mean
33 occurrence over the year ($\chi^2(3) = 81.13, p < .001$; 2-month: 25.70%, 4-month: 13.50%,
34
35 6-month: 9.85% and 12-month: 3.56%). Post-hoc tests revealed significant decreases in
36
37 the occurrence of Blue facial expressions over age, with 2-month old infants displaying
38
39 significantly more Blue expressions than 4-, 6-, and 12-month olds and 12-month olds
40
41 displaying significantly fewer Blue expressions than 4- and 6-month olds. At all ages,
42
43 infants displayed the most Blue facial expressions within the first 5 seconds after the
44
45 immunization needle and decreased steadily within the first 15 seconds. Consistently over
46
47 the minute, infants at 2 months displayed the most Blue facial expressions, followed by
48
49 infants at 4, 6, and 12 months. However, over the last 40 seconds, the 4-, 6- and 12-
50
51 month trajectories appeared indistinguishable.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Purple: There were no age differences in average occurrence of Purple facial expressions ($\chi^2(3) = 5.83, p = 0.12$; 2-month: 1.40%, 4-month: 1.76%, 6-month: 1.87% and 12-month: 1.93%). The amount of Purple facial expressions remained low and flat across the post-immunization minute for each age. Overall, infants consistently displayed Purple facial expressions less than 2% of the time after the immunization needle.

Black (see Figure 6): Black facial expressions significantly differed in mean occurrence across age ($\chi^2(3) = 25.42, p < .001$; 2-month: 4.26%, 4-month: 3.20%, 6-month: 3.60% and 12-month: 11.24%). Post hoc pairwise comparisons revealed significant increases in Black facial expressions from 2, 4, and 6 months to 12 months. Lastly, for each age Black facial expressions had a slight tendency to increase over the one minute following the immunization needle. Infants at 12 months displayed consistently more Black facial expressions across the entire minute compared to the younger ages.

Discussion

Infants' facial actions were comprehensively coded following an acutely painful procedure until one minute after a noxious stimulus (i.e., immunization). Many of the categories of facial expressions empirically identified in this study involved some of the same components as the adult pain expressions (i.e., Red, Orange, Yellow, Green, and Blue expressions). In fact, most of the variation in adult facial expressions in response to pain can be limited to four facial actions [32]: brow lowering (AU4), cheek raising/eye lid tightening (AU6/AU7), nose wrinkling/lip raising (AU9/AU10), and eye closure (AU43). As with the adult facial expression of pain [32], the degree of vertical mouth

opening (AU26/AU27) was not a sensitive or specific component of the infant facial expression of pain. However, a distinct difference emerged between expressions of pain in adults and infants. Specifically, a horizontal stretching of the mouth (AU20) has not always been associated with expressions of pain in adults [32,35], but this study suggests that it is an important component of the infant facial expression of pain.

Horizontal stretching of the mouth is likely more integral to infant facial expressions of pain because, along with mouth opening, it is the **principal** component of cry faces and is usually associated with cry vocalizations. Adults have more control over their pain responses and rarely cry in response to an acute pain stimulus such as an immunization. Most infants, however, cry in response to those very same stimuli. More generally, crying is the first form of vocal communication exhibited by infants after birth and the most common response to generalized distress or discomfort. Therefore, the intensity of horizontal stretching of the mouth may be considered when assessing pain intensity in infant facial expressions. This idea is supported by work in the infant pain literature [14,15].

In this study we examined the facial expressions of pain among infants at four different ages in a systematic fashion. This study went further than earlier studies because we coded the entire spectrum of facial actions, not just those presumed to be associated with pain. NFCS [14,15], a measure of pain, has been shown to lack the ability to differentiate between pain-related and non-pain-related distress [1]. One challenge in defining differing emotional expressions, regardless of age, is the overlap of facial actions involved in all negative emotional expressions. Past studies with infants [3,19,20]

1
2
3
4
5
6 ignored this commonality in their attempts to code discrete negative emotions in infants'
7
8 distress responses. However, in pain contexts this overlap makes tremendous sense. An
9
10 influential review regarding the affective dimension of pain [31] describes a complex,
11
12 simultaneous affective reaction to pain rather than a unitary response. The findings herein
13
14 demonstrate that in the first year of life, there is a core set of facial actions in response to
15
16 pain and regulation of distress from pain. This study has further established that this core
17
18 set has seven readily discernible constellations. We speculate that, owing to a common
19
20 core set of facial actions, these seven constellations do not reflect discrete negative
21
22 emotions but rather different variants and intensities of sensory or affective pain
23
24 responses to noxious stimuli.
25
26
27
28
29

30
31 Although we coded every visible facial muscle action using a fine-grained,
32
33 anatomically based coding system, we did not observe expressions of discrete negative
34
35 emotions in the immediate post-needle epoch. Table 3 illustrates the facial expressions
36
37 identified in this study and their suggested interpretations according to the BabyFACS
38
39 [23], NFCS [14,15], and Max [17] manuals as well as an intensity metric for adult pain
40
41 [32] specified in terms of FACS AUs [10,11]. As seen in Table 3, the constellations of
42
43 facial expressions shown by infants immediately following immunization do not
44
45 correspond to prototypical, adult-like expressions of discrete negative emotions. Instead,
46
47 Table 3 demonstrates that according to NFCS, BabyFACS, and the adult pain metric, all
48
49 faces would be classified as distress or pain. However, according to the Max system, only
50
51 Blue would be classified as discomfort-pain. Yellow would be coded as discrete anger,
52
53 and other configurations would be coded as blends of different negative emotions. It is
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6 important to note that the Max formulas for discrete negative emotions are simply based
7
8 on how widely open an infant's eyes and mouth are. While it is feasible to suggest that
9
10 the degree of mouth and eye openness reflects discrete emotions, these distinctions have
11
12 not been supported by empirical research [7,24,25]. We suggest that these variations in
13
14 the intensities of distress configurations may reflect sensory intensity differences and
15
16 developing abilities to regulate distress.
17
18
19

20
21 Earlier, we suggested that the seven constellations identified here reflect different
22
23 pain variants. From what is generally known about the immunization experience and
24
25 acute procedural pain, pain is felt the most intensely initially (i.e. from the skin breakage
26
27 and the vaccine being pushed into the skin or muscle) and gradually abates as a function
28
29 of time [27,30]. Thus, it appears logical that facial expressions closest in time to the
30
31 needle may be more representative of the most intense pain expression (high noxious
32
33 sensory stimulation with initial reactivity) while facial expressions closer to one minute
34
35 post-needle may reflect blended expressions of negative emotions and less intense
36
37 sensory pain as well as efforts to regulate pain and distress.
38
39
40
41

42
43 Our findings are consistent with research demonstrating that infants do not show
44
45 differentiated facial expressions of discrete negative emotions like sadness and fear in the
46
47 first year of life [7,24] in response to pain. It also shows that there is not a single,
48
49 stereotyped pain expression during infancy. Infant facial expressions serve the purpose of
50
51 non-verbally communicating a need or a desire [28,29]. When young infants experience
52
53 pain (and negative emotion associated with pain), they need to communicate that they
54
55 need help. A generalized distress expression serves a strong evolutionary function in that
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6 it would likely result in proximity from a caregiver [21]. As gaining proximity or help
7
8 from a caregiver quickly is the likely end goal for infant pain communication, **it seems**
9
10 **crucial** for a caregiver to discern exactly why the young infant needs help and more
11
12 efficient to simply be signaled that help is needed urgently and that proximity is
13
14 necessary. Consistent with previous ethological thinking and an ontogenetic perspective,
15
16 we agree that a generalized infant cry is enough to engage the caregiver system regardless
17
18 of the specific cause [21].
19
20
21

22
23 By tracking the occurrence of infants' facial expressions over the first minute
24
25 post-needle and the decrease of acute pain responding in infancy over the first year of
26
27 **life, we** can attempt to speculate about the communicative goals of the facial expressions
28
29 discerned and how they may change as the infant ages. The goal of this study was to
30
31 provide the field with a broad inventory of all possible infant facial muscle configurations
32
33 that occur in an acute pain context. We aimed to gain novel data germane to
34
35 understanding the development and regulation of negative affect over the first year of
36
37 life. Our analysis of the temporal trajectories of the seven categories of facial responses
38
39 we identified suggested that these are pain related distress expressions ranging from the
40
41 most intense pain and sensory overload immediately following inoculation (Blue
42
43 expressions with tightly closed eyes: "complete sensory overwhelm") to decreasing
44
45 degrees of pain (Green, with closed eyes and less intense cry mouth: "moderate sensory
46
47 overwhelm") and increasing capacity with time and age to regulate distress (Red, Orange,
48
49 and Yellow expressions). The Yellow expressions, with a big cry mouth and open eyes,
50
51 may reflect attempts to regulate intense distress and maintain visual contact ("regulation
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6 from intense distress”); Red expressions may be labeled “regulation from moderate
7
8 distress”; and Orange expressions may be labeled “regulation from mild/moderate
9
10 distress”. Lastly, facial configurations with less intense cry face components might be
11
12 labeled “mild distress” (Purple) or “mild discomfort” (Black). We propose that these
13
14 seven categories of expressions may have evolved to allow infants to communicate
15
16 crucial two broad states to caregivers: level of distress and degree of regulation from
17
18 distress. Although these categories of facial expressions are visually distinguishable,
19
20 within the context of pain, we suggest that they do not represent expressions of discrete
21
22 negative emotions.
23
24
25
26

27 *Study Limitations*

28
29
30 Infants did not receive the same immunization at 12 months as they did at 2, 4,
31
32 and 6 months [36]. However, regardless of vaccine protocol, all infants were exposed to a
33
34 highly painful acute stimulus and showed modulation of the intensity of distress
35
36 expressions over the minute. Nevertheless, following infants longitudinally at 2, 4, 6 and
37
38 12 months provided evidence of developmental changes in the relative frequency of the
39
40 different expressions observed and their time course.
41
42
43

44
45 Our analysis of expressions was limited to facial expressions. Caregivers almost
46
47 always have access to vocalizations and body movements when judging an infant’s
48
49 emotions; therefore, there is strong merit in considering these expressive modalities in
50
51 relation to facial expressions. However, facial expressions collectively are used by
52
53 infants, children, and adults to communicate the infant’s emotions and behavioral
54
55 dispositions to others [4,9,24] and are therefore are critical in their own right.
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6 Since our analyses focused on responses to pain within the first minute post
7
8 needle, we cannot draw any general conclusions outside this specific context. However,
9
10 our findings are consistent with those of studies that have emphasized the importance of
11
12 infants' facial expressions for parents' ability to read and respond sensitively to their
13
14 infants' distress [21,24,27].
15
16

17 18 **Acknowledgements**

19
20 This research was supported by funds from the Canadian Institutes of Health Research
21
22 (New Investigator Award to Dr. Pillai Riddell, Operating Grant [PI: Dr. Pillai Riddell]),
23
24 the Ontario Graduate Scholarship Program (Doctoral Award to Ms. Ahola Kohut) and the
25
26 Social Sciences and Humanities Research Council (Doctoral Award to Ms. Ahola
27
28 Kohut). The authors thank the staff and families at our research clinics for their assistance
29
30 and ongoing support as well as the specific infants and caregivers who gave permission to
31
32 publish their photos.
33
34
35

36 37 **Conflict of Interest Statement**

38
39 The authors have no financial interest in the results of this research. None of the authors
40
41 have any conflict of interest with this work.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- [1] Ahola Kohut S, Pillai Riddell RR. Does the NFCS discriminate between infants experiencing pain related and non-pain related distress? *J Pain* 2009;10:214-220.
- [2] Anand KJ. Effects of perinatal pain and stress. *Prog Brain Res* 2000;122:117-129.
- [3] Axia G, Bonichini S, Benini F. Attention and reaction to distress in infancy: A longitudinal study. *Dev Psychol* 1999;35:500-504.
- [4] Bowlby J. *Attachment and loss: vol.1. Attachment*. New York: Basic Books; 1969/1982.
- [5] Bradley S. *Affect Regulation and the Development of Psychopathology*. New York: Guilford Press; 2000.
- [6] Bridges KM. Emotional development in early infancy. *Child Dev* 1932;3:324-341.
- [7] Camras LA, Oster H, Bakeman R, Meng Z, Ujiie T, Campos JJ. Do infants show distinct negative facial expressions for fear and anger? Emotional expression in 11-month-old European, American, Chinese, and Japanese infants. *Infancy* 2007;11:131-155.
- [8] Camras LA, Witherington DC. Dynamical systems approaches to emotional development. *Dev Rev* 2005;25:328-350.
- [9] Darwin C. *The Expression of the Emotions in Man and Animal*. Great Britain: John Murray Press; 1872.

1
2
3
4
5
6 [10] Ekman P, Friesen WV. Measuring facial movement. *J Nonverbal Behav* 1976;1:56-
7
8 75.
9

10 [11] Ekman P, Friesen WV, Hager J. Facial Action Coding System. [CD ROM] Salt Lake
11
12 City: Research Nexus, Subsidiary of Network Information Research Corporation; 2002.

13 [12] Fernald CD, Corry JJ. Empathic versus directive preparation of children for needles.
14
15 *Child Health Care* 1981;10:44-47.
16
17

18 [13] Fitzgerald M. The development of nociceptive circuits. *Nature Rev Neurosci*
19
20 2005;6:507-520.
21
22

23 [14] Grunau RV, Craig KD. Neonatal Facial Coding System: Revised Training Manual.
24
25 Vancouver: Univ of British Columbia; 2010.
26
27

28 [15] Grunau RV, Craig KD. Pain expression in neonates: facial action and cry. *Pain*
29
30 1987;28:395-410.
31
32

33 [16] Izard CE. On the ontogenesis of emotions and emotion-cognition relationships in
34
35 infancy. In: Lewis M, Rosenblum LA. editors. *The Development of Affect* . New York:
36
37 Plenum; 1978. p.389-413.
38
39

40 [17] Izard CE. The maximally discriminative facial movement coding system (Max).
41
42 Newark: Univ. of Delaware; 1979 (Revised, 1995).
43
44

45 [18] Izard CE. *The psychology of emotions*. New York: Plenum Press; 1991.
46
47

48 [19] Izard CE, Hembree EA, Dougherty LM, Spizzirri CL. Changes in facial expressions
49
50 of 2- to 19-month-old infants following acute pain. *Dev Psychol* 1983;19:418-426.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6 [20] Izard CE, Hembree EA, Huebner RR. Infants' emotion expressions to acute pain:
7
8 Developmental change and stability of individual differences. *Dev Psychol* 1987;23:105-
9
10 113.
11
12 [21] Mesman J, Oster H, Camras L. Parental sensitivity to infant distress: What do
13
14 discrete negative emotions have to do with it? *Attach Hum Dev* 2012; in press.
15
16 [22] Messinger DS. Positive and negative: Infant facial expressions and emotions. *Curr*
17
18 *Dir Psychol Sci* 2002;11:1-6.
19
20 [23] Oster H. *Baby FACS: Facial Action Coding System for Infants and Young Children*.
21
22 New York: New York Univ.; 2010.
23
24 [24] Oster H. The repertoire of infant facial expressions: An Ontogenetic perspective. In:
25
26 Nadel J, Muir D. editors. *Emotional development: Recent Research Advances*. New
27
28 York: Oxford Univ.; 2005. p. 261-292.
29
30 [25] Oster H, Hegley D, Nagel L. Adult judgments and fine-grained analysis of infant
31
32 facial expressions: Testing the validity of a priori coding formulas. *Dev Psychol*
33
34 1992;28:1115-1131.
35
36 [26] Peters JW, Schouw R, Anand KJ, van Dijk M, Duivenvoorden HJ, Tibboel D. Does
37
38 neonatal surgery lead to increased pain sensitivity in later childhood? *Pain* 2005;114:444
39
40 – 454.
41
42 [27] Pillai Riddell R, Campbell L, Flora D, Racine NM, Din Osmun L, Garfield H,
43
44 Greenberg S. The relationship between caregiver sensitivity and infant pain behaviors
45
46 across the first year of life. *Pain* 2011;152:2819-2826
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6 [28] Pillai Riddell RR, Chambers CT. Parenting the infant in pain. In: Anand KJS, B.
7
8 Stevens B, McGrath PJ, editors. Pain in neonates and infants. Edinburgh: Elsevier Ltd.;
9
10 2007. p. 279-289.
11
12
13 [29] Pillai Riddell RR, Racine NM, Assessing pain in infancy: The caregiver context.
14
15 Pain Res Manag 2009;14:27-32
16
17
18 [30] Pillai Riddell RR, Stevens B, Cohen LL, Flora D, Greenberg S. Predicting maternal
19
20 and behavioural measures of infant pain: The relative contribution of maternal factors.
21
22 Pain 2007;133:138-149.
23
24
25 [31] Price DD. Psychological and neural mechanisms of the affective dimension of pain.
26
27 Science 2000;288:1769-1772.
28
29
30 [32] Prkachin KM. The consistency of facial expressions of pain: A comparison across
31
32 modalities. Pain 1992;51:297-306.
33
34
35 [33] Prkachin KM. Assessing pain by facial expression: Facial expression as nexus. Pain
36
37 Res Manag 2009;14:53-58.
38
39
40 [34] Prkachin KM, Craig. Expressing pain: The communication and interpretation of
41
42 facial pain signals. J Nonverbal Behav 1995;19:191-205.
43
44
45 [35] Prkachin KM, Mercer SR. Pain expression in patients with shoulder pathology:
46
47 Validity, properties and relationship to sickness impact. Pain 1989;39:257-265.
48
49
50 [36] Schechter N, Zempsky WT, Cohen LL, McGrath PJ, McMurtry CM, Bright NS.
51
52 Pain reduction during pediatric immunizations: Evidence-based review and
53
54 recommendations. Pediatrics 2007;119:1184-1198.
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

[37] Sroufe LA. Emotional Development: The Organization of Emotional Life in the Early Years. New York: Cambridge University Press; 1996.

Figure Captions

Figure 1. Simplified schematic representation of the categorization of facial expression.

Figure 2. Example of a Red facial expression and graph presenting the proportion of Red facial expressions displayed in response to last needle.

Figure 3. Example of a Yellow facial expression and graph presenting the proportion of Yellow facial expressions displayed in response to last needle.

Figure 4. Example of a Green facial expression and graph presenting the proportion of Green facial expressions displayed in response to last needle.

Figure 5. Example of a Blue facial expression and graph presenting the proportion of Blue facial expressions displayed in response to last needle.

Figure 6. Example of a Black facial expression and graph presenting the proportion of Black facial expressions displayed in response to last needle.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Summary Statement

Longitudinal analysis of facial expressions suggests that child/adult negative emotion prototypes are not seen in infants but rather pediatric variants of pain/distress expressions.

Figure 1




Figure 1. Simplified schematic representation of the categorization of facial expression.

Figure 2




Figure 2. Example of a Red facial expression and graph presenting the proportion of Red facial expressions displayed in response to last needle.




Figure 3. Example of a Yellow facial expression and graph presenting the proportion of Yellow facial expressions displayed in response to last needle.




Figure 4. Example of a Green facial expression and graph presenting the proportion of Green facial expressions displayed in response to last needle.




Figure 5. Example of a Blue facial expression and graph presenting the proportion of Blue facial expressions displayed in response to last needle.




Figure 6. Example of a Black facial expression and graph presenting the proportion of Black facial expressions displayed in response to last needle.

Table 1. Demographic Variables and Characteristics of Participants (N=100)

	Frequency	Percentage
Infant Sex (% male)	50	50.00%
Parent-reported Heritage Culture		
North American	17	17.00%
European	47	47.00%
Asian	11	11.00%
South/Latin American	1	1.00%
Middle Eastern/African	2	2.00%
Jewish	8	8.00%
Other	8	8.00%
Mixed Canadian	6	6.00%
Mother's highest education		
Graduate school/Professional training	30	30.00%
University graduate (4 years)	44	44.00%
Partial university (at least 1 year)	4	4.00%
Trade school/Community college	16	16.00%
High school Graduate	6	6.00%
Marital status		
Married	87	87.00%
Divorced/Separated	2	2.00%
Never Married	1	1.00%
Other	1	1.00%
Common Law	8	8.00%
Engaged	1	1.00%

Table 2. Descriptions of the Categories of Most Common Negative Facial Expressions

Categories	Red	Orange	Yellow	Green	Blue	Purple	Black
Distinguishing Characteristics ^a	Any cry mouth with oblique brows	Horizontal cry mouth with eyes open	Big cry mouth (horizontal & vertical) with eyes open	Horizontal cry mouth with eyes closed	Big cry mouth (horizontal & vertical) with eyes closed	Any cry mouth without cheek raise or eyelid tightening	Non-cry negative mouth
AUs							
1+3, 4	REQ	NO	NO	NO	NO	NO	OP
1+2+3, 4	NO	NO	OP	NO	OP	OP	NO
2+3, 2+4	NO	OP	OP	OP	OP	NO	OP
43	NO	NO	NO	REQ	REQ	OP	OP
6, 7	REQ	REQ	REQ	REQ	REQ	NO	OP
20	REQ	REQ	REQ	REQ	REQ	REQ	NO
25+26ab	OP	OP	NO	OP	NO	OP	OP
25+26c/27	OP	NO	REQ	NO	REQ	OP	OP
17	OP	OP	NO	OP	OP	OP	OP

Note. REQ = required, OP = optional, NO = not allowed, ^aDetailed verbal descriptions of categories can be found in Table 3

Table 3. Descriptions and Interpretations of the Categories of Negative Expressions

Category Name	Description of Expression	NFCS ^{a*}	MAX ^b	BabyFACS ^c	Adult Pain ^d
Red	<ul style="list-style-type: none"> - brow oblique and drawn together - cheek raise and/or eyelid tightening - lip raise or nasolabial furrow - slightly to moderately open mouth - horizontal stretch of the mouth 	- Pain score: 3-5	- Sadness/Anger blend - Sadness/Fear blend	- Pre-cry or cry face - Fuss face - Moderate distress	- Pain score ^d : 9-12
Orange	<ul style="list-style-type: none"> - brow knitting and/or knotting - cheek raise and/or eyelid tightening - lip raise or nasolabial furrow - slightly open mouth - horizontal stretch of the mouth 	- Pain score: 4-5	Anger/Fear Blend	- Grimace - Fuss face - Mild to moderate distress	- Pain score ^d : 9-12
Yellow	<ul style="list-style-type: none"> - brow knitting and/or knotting - cheek raise and/or eyelid tightening - lip raise or nasolabial furrow - widely open mouth - intense horizontal stretch of the mouth 	- Pain score: 5-6	- Anger	- Intense cry face - Intense distress	- Pain score ^d : 9-12
Green	<ul style="list-style-type: none"> - brow knitting and/or knotting - cheek raise and/or eyelid tightening - eyes closed - lip raise or nasolabial furrow - slightly open mouth - horizontal stretch of the mouth 	- Pain score: 5-6	- Discomfort-Pain/Fear blend	- Moderate cry face - Fuss face - Moderate distress	- Pain score ^d : 10-13
Blue	<ul style="list-style-type: none"> - brow knitting and/or knotting - cheek raise and/or eyelid tightening - eyes closed - lip raise or nasolabial furrow - widely open mouth - intense horizontal stretch of the mouth 	- Pain score: 6-7	- Discomfort-Pain	- Intense cry face - Intense distress	- Pain score ^d : 10-13

Purple	- brow knitting and/or knotting - lip raise or nasolabial furrow - slightly to moderately open mouth - horizontal stretch of the mouth	- Pain score: 3	- Partial anger	- Mild cry face - Fussy	- Pain score ^d : 6-8
Black	- brow knitting and/or knotting - lip raise or nasolabial furrow - lip corners may be depressed - possible pout or “horseshoe” mouth	- Pain score: 2-3	- Anger/Sadness blend - Discomfort- Pain/Sadness blend	- Pouts - Horseshoe-mouth - Emotion regulation - Modulated distress	- Pain score ^d : 6-8

Note: ^a NFCS Manual [14,15], ^b MAX Manual [17], ^c BabyFACS Manual [23], ^d Prkachin’s [32,35] adult pain intensity metric based on FACS Action Units [10,11]